

Annals of the Missouri Botanical Garden

Vol. 10

FEBRUARY, 1923

No. 1

POTATO BLACKLEG WITH SPECIAL REFERENCE TO THE ETIOLOGICAL AGENT¹

HARRY MILLIKEN JENNISON

*Professor of Botany, University of Tennessee
Formerly Assistant in Botany, Henry Shaw School of Botany of
Washington University*

INTRODUCTION

The objects of the investigations presented herewith were several, but only those outstanding should be mentioned at this point. First of all, a determined effort was made to discover the relationships, one to another, of the several "species" of bacteria recorded as being the cause of the blackleg disease of potatoes. To this end a thorough comparative study of the morphology, cultural features, and physiology of some 12 strains of the blackleg bacillus was made. The cultures employed represented strains of the organism from regions widely separated geographically. Among them were the 4 "species" originally described as being the cause of the disease in question. In the prosecution of the problems arising the writer was led into a quantitative study of carbohydrate utilization by strains of the blackleg bacillus and other microorganisms. The work done in this connection constitutes an important phase of the investigations carried out. By way of extending the usefulness of the paper, the writer presents a rather full and complete diagnosis of the disease, with a discussion of the economic aspects, also a revised description of the potato blackleg parasite.

¹An investigation carried out at the Missouri Botanical Garden in the Graduate Laboratory of the Henry Shaw School of Botany of Washington University, and submitted as a thesis in partial fulfillment for the requirements for the degree of doctor of philosophy in the Henry Shaw School of Botany of Washington University.

HISTORICAL

Frank ('97, '99) was the first to publish accurate and detailed descriptions of the potato blackleg disease. One may be assured he was discussing the potato disease which goes by the name of "blackleg" and which is so common and well known in many parts of North America. He reported the cause of the disease to be a bacterium which he briefly described and named *Micrococcus phytophthorus*. Micrococci have been frequently met with in isolation plates made from potato stems and tubers affected with the blackleg disease. Many other workers have reported similar experiences, but no one has presented adequate proof that any one of the species of *Micrococcus* found present under such circumstances is pathogenic to the potato. Until such time as a species of *Micrococcus* conforming to Frank's description shall be proved capable of infecting potatoes and causing the blackleg disease, *Micrococcus phytophthorus* Frank should be considered a *nomen nudum*.

About 20 years earlier Hallier ('78) described an infectious wet rot of the potato and he discussed at considerable length the association of certain bacteria ("Vibrionen," "Bakterien," "Micrococcus") and a fungus (*Peronospora infestans* Casp.) in the affected potatoes. One cannot determine from the text which of the organisms is of primary significance, but as a mere matter of historic interest it may be noted that this writer contended that the wet rot described by him was caused by "die Produkte der Plastiden der *Peronospora infestans* Casp." While it appears unlikely that the wet rot described by Hallier is in any way related to the rot due to the blackleg germ, it is nevertheless interesting to note the mention at this early date of a potato tuber rot in which bacteria were found present.

D'Arbois de Jubainville and Vesque ('78), nearly 45 years ago, showed that they were familiar with a potato tuber rot ("pourriture cellulaire") occurring in the soil, but the bacteria were not identified as the cause; rather, the disease was thought by them to be due to an excess of nutrients and water.

A year later Reinke and Berthold ('79), as a result of studies of potato decay due to fungi, concluded that a wet rot of potato tubers existed which was not due to fungi. Furthermore, they demonstrated the association of bacteria with the wet rot described, and they also proved the infectious nature of the rot.

Burrill ('90, '93) published brief notes on a bacterial disease of potatoes, but as nearly as can be determined, the disease which he studied was the brown rot of potato, later most thoroughly investigated by E. F. Smith.

A bacterial malady of the potato was described by Prillieux and Delacroix ('90) in France, and what they took to be the causal organism was designated *Bacillus caulivorus*. The potato disease referred to by Prillieux and Delacroix was described as one which affects the stem, beginning at the bottom and working upwards. To this extent, at any rate, it is comparable to the one under consideration. No description of the cultural characteristics of the organism was presented in their paper and no statement was made concerning the method of isolation by which the organism was obtained. A few years later Prillieux ('95) briefly mentioned certain of its characteristics, stating that it developed a green coloration in some media. Laurent ('99) stated that *B. caulivorus* was probably the common saprophyte *B. fluorescens liquefaciens* Flügge. Later, Delacroix concluded, according to Prunet, that *B. caulivorus* was most probably *B. fluorescens liquefaciens* Flügge, which he thought became parasitic under certain conditions. A more complete review of the work referred to just above will be found in Prunet's ('02) paper.

One of the early contributions to our knowledge of bacterial diseases of plants in general and of the potato in particular is that of Kramer ('91), who describes quite fully a wet rotting of potatoes. He obtained pure cultures of the supposed etiological agent which he described as a spore-bearing, rod-shaped bacterium.

The potato disease described by Tyron ('94, '99) as occurring in Queensland, Australia, is probably not the blackleg disease. Rather according to Tyron's statements, it is very similar to, if not the same as, E. F. Smith's brown rot disease (see especially Tyron, '99, "Biography," p. 62, and footnote p. 63; also Smith, '14, p. 208).

Smith's first publication on the brown rot of *Solanaceae* caused by *Pseudomonas* (*Bacterium*) *solanacearum* E. F. S. appeared in 1896. From the outset there was little or no confusion of this disease with the one under discussion. This paper by Smith is one of the first thoroughgoing and accurate descriptions of a bacterial disease of the potato. Recalling that his work was

done at an early date, and at a time when foreign bacteriologists of high standing opposed the idea that bacteria could be the primary cause of disease in plants, the contribution is all the more significant.

Wehmer ('98) published extensively the results of his investigations of potato diseases. In Part 3 he describes a bacterial rot of the tuber, but he definitely takes the attitude that the bacteria found associated with the rot were not the primary etiological agents. His attitude may be taken as representative of the belief of most German botanists and bacteriologists of that time.

There occurred in the vicinity of St. Petersburg, Russia, in 1898, a potato bacteriosis which was described by Iwanoff ('99). The causal organism was described as a short, oval-cylindrical, active rod found (in the tissues) to measure $1.5\mu \times 0.5\mu$. The disease was one which affected the leaves and stems, early symptoms being manifest in the leaves. Later, the stems were affected and showed symptoms of the disease. Brown lesions appeared externally on the stems, the pith was attacked and destroyed, and the stems wilted and died. Starch was not destroyed. Iwanoff contended that the disease he described was similar to that described by Smith ('96) as due to *B. solanacearum*. It appears to the writer that he was mistaken in the view presented, and from our limited knowledge of the situation it seems more likely that the potato disease prevalent in Russia near St. Petersburg in 1898 was the blackleg disease.

Jensen ('00) investigated a bacterial disease of the potato which he referred to as "blackleg". As a result of microscopic investigations of potato plants affected with "blackleg" he concluded that the disease was caused by bacteria ("Mikrokokken").

What Smith ('14) refers to as "The French Disease" was first described by Delacroix ('01) in a succession of short papers appearing in 1901. Delacroix first stated that the potato malady due to *B. solanacearum* E. F. Smith was prevalent in France. Later, he concluded that the disease referred to above, the one for which he suggested the name "brunissure," was caused by a bacterium new to science which he named *Bacillus solanicola*. Smith ('14), however, claimed that, "Here again, it is uncertain whether we have to do with *Bacillus phytophthorus*, *Bacterium solanacearum*, or some third organism. The writer obtained a

culture of *B. solanicola* from Prof. L. R. Jones, to whom it was given in Paris by Delacroix, but either it never possessed any pathogenic properties, which is quite probable, or else had lost them by cultivation." The same investigator made the following statement, after having had opportunity to examine material collected and preserved by Delacroix to illustrate the disease: "It represents a fungous disease of the potato."

Part III of van Hall's ('02) doctoral dissertation ("The stem rot or blackleggedness of potato stems caused by *Bacillus atrosep-ticus* nov. sp.") gives a fairly complete description of the disease under discussion, together with a diagnosis of the etiological agent sufficiently detailed and complete to enable one to classify the parasite. For reasons which appear below, the writer is of the opinion that the potato disease commonly referred to throughout the United States and many parts of Europe as "blackleg" is one and the same thing as that described by van Hall. The etiological agent in all cases is probably identical and, as far as is known, should be referred to van Hall's *Bacillus atrosep-ticus*.

At a slightly earlier date Appel ('02) published a short paper ("Contribution to our knowledge of the bacterial rotting of potatoes"), in which he described effects of the rot, also experiments on the pathogenicity of the disease and stated that the systematic position of the causal agent of the potato rot, as well as its practical importance, would be given in a forthcoming article. About a month later another short paper was published by the same investigator (Appel, '02a), entitled "The cause of the 'Blackleggedness' of the potato." In this article he takes exception to Frank's binomial in the following words: "Wenn nun auch der Frank'sche Name *Micrococcus phytophthorus* auf den *Bacillus* nicht anwendbar ist, so glaube ich doch aus praktischen Gründen den Speciesnamen beibehalten zu sollen, um so mehr als es sich herausstellte, dass eine ganze Reihe von Pflanzen in charakteristischer Weise angegriffen wird. Ich nenne daher den von mir isolirten *Bacillus*, welcher Schwarzbeinigkeit und Knollenfaule bei den Kartoffeln hervorruft: *Bacillus phytophthorus* Appel." Nowhere in this paper, however, is there to be found a description of the morphological and cultural characters (other than those implied by the name) of the etiological agent which he had isolated. The binomial given by Appel in this publication must therefore be regarded as a *nomen nudum*.

In 1903 Appel ('03) published a complete account of the potato blackleg disease as it occurs in Germany, together with a description of the causal organism. The same investigator (Appel, '05, '06) contributed substantial additions to our knowledge of this disease.

Butler ('03) mentions a bacterial disease which he thought was possibly the same as that described in the United States by E. F. Smith under the name of brown rot.

Jones ('05), in an account of disease resistance of potatoes, presented a description of the blackleg disease and accompanied it with observations upon its occurrence in Europe. He reported that it was found in Holland, Belgium, Germany, France, and England. The first authentic record of the occurrence of potato blackleg in the United States is that of Jones ('07) who found it in 1906 on a farm in Vermont. The seed used in planting the field came from Maine. He states also: "Vague reports have frequently come to the Station in previous years as to troubles of this class . . ." This author (Jones, '07) "passed almost directly from German fields where it was prevalent, to English fields and found the malady equally common and identical in appearance with that on the continent." This statement is of especial significance in connection with the report of Johnson ('07) who claims to have gathered substantial evidence of the existence of the potato blackleg disease in Ireland. (Compare also with the report made by Pethybridge and Murphy in 1910).

Harrison ('07) published a very complete account of a potato rot occurring in eastern Canada. The symptoms of the disease described by him were, he states, quite similar to the "Schwarzbeinigkeit" of van Hall and Appel. While he attributes the disease to a bacterial species (*Bacillus solanisaprus*) new to science, the writer has reason to believe (see below) that the causal agent described by him is identical with *Bacillus atrosep-ticus* van Hall. Morse ('07) found potatoes affected with blackleg in Maine in 1907. In 1909 he (Morse, '09, '11) published an account of the nature of the disease, its distribution, economic importance, and its control in Maine.

Smith ('10), however, was the first American to publish a full account of the blackleg parasite. His description is based on studies made with a subculture of Appel's species which he obtained from Aderhold in Berlin, presumably a transfer from

Appel's original culture. Smith states in this article that he regards *Bacillus solanisaprus* Harrison as closely related to, but not identical with, *B. phytophthorus* Appel.

About the same time Pethybridge and Murphy ('10) published a note on a bacterial disease of the potato plant, attributing it to a new species of bacterium for which they proposed the name "*Bacillus melanogenes*." In 1911 these investigators published a full account of the malady referred to above, together with a complete diagnosis of the etiological agent. The potato disease described by them was probably none other than the "blackleg," and there is very little, if any, doubt in the writer's mind that their *Bacillus melanogenes* should be referred to *Bacillus atrosepticus* van Hall.

Murphy ('16) gives an account of a blackleg disease of potatoes occurring in Canada, attributing the disease to *Bacillus melanogenes* Peth. & Murphy.

From a series of observations made in 1916, Morse ('17) reports having seen potato blackleg in certain of our western states, though he found it to differ in certain respects from that familiar to him in Maine. This paper is an important contribution to the literature of the subject. Of particular significance are his comparative studies of the causal organisms. The writer is in accord with Morse's conclusion that the strains studied (including cultures "received under the names '*Bacillus atrosepticus* van Hall,' '*B. solanisaprus* Harrison,' and '*B. melanogenes* Peth. & Murphy,'" as well as 3 isolated from blackleg material in Maine) are identical. However, since Morse was unable to procure a trustworthy culture of *B. phytophthorus* Appel he presents no data which "bear on the relationship between the organism originally described by Appel as *B. phytophthorus* and the other strains of blackleg bacteria." So in deference to the opinion of Smith ('10) that *B. phytophthorus* Appel is not identical with *B. solanisaprus* Har., Morse was forced to exclude it from *B. atrosepticus* van Hall. This is most regrettable, since Morse shows that small differences in the physiology of bacteria are not sufficient for the establishment of new species. His work shows abundant evidence of having been carefully done. Unfortunately, there seems to be no record of the comparative studies upon which Smith ('10) bases his opinion.

Paine ('17) referred to the cause of the potato blackleg studied by him in England as *Bacillus atrosepticus* van Hall and ex-

pressed the opinion that the species described by Appel, Harrison, and Pethybridge and Murphy should be referred to van Hall's *Bacillus atrosepticus*. The argument advanced by this investigator is most logical, but attention is called to the fact that he did not carry out comparative studies using the several "species" in question.

Rosenbaum and Ramsey ('18) contributed the results of some studies upon the influence of temperature and precipitation on the blackleg of potato.

Ramsey's ('19) studies of the viability of the potato blackleg organism led him to believe that the pathogen does not live in the tubers left in the field over winter. He expressed the further conclusion that unless the potato seed were infected at planting time there was little chance that the uninjured plants would contract the disease.

Artschwager's ('20) researches upon the pathological anatomy of potato blackleg form a new contribution to our general knowledge of this disease. He studied only blackleg plants grown in the arid regions of western Colorado. The affected plants examined by him were found to show an increase in strongly lignified vascular tissue and a transformation of part or most of the parenchyma cells of the cortex and pith into sclerids. This investigator also discovered that protein crystals occurred in great abundance in all organs of plants affected with the blackleg disease, especially in the leaves, while under normal conditions protein crystals have been observed only in the peripheral cell layers of the cortex of the potato tubers.

Jennison's ('21) abstract, "*Bacillus atrosepticus* van Hall, the cause of the blackleg disease of Irish potatoes" was based on the investigations carried out in large part in 1916 and 1917, here reported upon in detail. While the group number as published by the writer at that time differs in respect to the last 3 digits from that of Morse ('17), it agrees throughout with that assigned by Shapovalov and Edson ('21). The last-named investigators contributed one of the most important of the recent accounts of the disease under discussion. They showed that the potato blackleg disease prevailed in the irrigated districts of the West, and that the causal agent was a *Bacillus*, which they concluded was "identical with *Bacillus phytophthorus* Appel in all the essential characteristics considered in determining bacterial spe-

cies." A detailed description of the causal organism isolated by them was given.

I. DIAGNOSIS OF THE DISEASE

As indicated in the preceding section, the blackleg disease of potatoes is a common and destructive bacterial disease of many varieties of *Solanum tuberosum*. The common name "blackleg" has been widely used in the United States and Canada, since it was first introduced by Jones in 1905. The term "blackleg" is a rather free translation of "Schwarzbeinigkeit," under which name the disease is known in Europe. The writer feels that blackstem and rot would be more suitable for American use but sees no point in suggesting that a change be made at this time, because (1) the old name has become well established and (2) because of its adoption by the American Phytopathological Society's Committee on Common Names for Plant Diseases. It may be pointed out, however, that the writer has observed that the term "blackleg" leads some farmers and others, especially in the West, to think that it may be related to the blackleg disease of cattle. More particularly is this the case when they learn that both are caused by bacteria.

Type—Potato blackleg is primarily a parenchyma necrosis, the cortical and pith tissues of both stem and tuber being almost exclusively involved. Occasionally, the vascular elements are found to be invaded by the parasite and sometimes they are somewhat browned.

Signs of blackleg in the field.—Considerable importance attaches to general manifestations of the disease as it appears in the field, since upon such manifestations depends the important practice of roguing, advocated below. Not infrequently "skips" or "missed hills" are due to this disease, for under favorable circumstances the young sprouts are destroyed by the parasite before they appear above ground; also the seed piece may be destroyed before it sprouts. When the disease progresses rapidly the plants wilt and become blackened to a considerable height. Such plants are quickly prostrated and soon die (pl. 1, fig. 3).

Secondary symptoms of the disease are likely to be the first to appear. Undersized vines with more or less yellowed and rolled leaves should be regarded with suspicion and more closely examined for the presence of black lesions upon the stem, both above

and below ground. There is, furthermore, a noticeable tendency for the upper leaves in particular to manifest a somewhat xerophytic texture and a starkness of growth-habit, the total effect being to give the tops of affected plants a narrowed or contracted appearance. Besides being light-colored, the upper leaves are often rendered more conspicuous by the presence of a metallic luster. Plants affected with blackleg, as a rule, offer little resistance to removal from the soil, and not infrequently break off when one attempts to pull them. If upon removal from the earth dark-colored cortical lesions are found on the underground portions of the stem, there is little doubt but that the plant is attacked by the blackleg organism. When upon splitting the stem longitudinally the pith is found blackened and more or less disintegrated the case is quite clear (pl. 1, figs. 2 and 4). Further confirmation is afforded if the seed piece has disintegrated. Aerial tuber development due to attack by the blackleg parasite is not commonly seen. The production of aerial tubers does take place when practically all the tuber-bearing stolons on well-developed plants are destroyed by the pathogen, thus precluding normal tuber development. *Rhizoctonia* more frequently causes such abnormalities of tuber development, and the underground lesions due to this fungus may be confused with those due to *Bacillus atrosepticus*. Certain characteristics of the *Fusarium* wilt disease may also be confused with the yellowing and wilting of potato vines caused by the blackleg parasite.

As a rule, plants affected with blackleg occur scattered promiscuously over the field. The occurrence and spread of the disease appear to bear a definite relation to environmental conditions, and the number of diseased plants is likely to be greater during a cold wet spring than in a warm dry one. Not infrequently there are more diseased plants in the low, poorly drained spots in a field.

"Vine" Symptoms.—The primary symptoms of the disease as they occur on the potato vines are most striking. These appear first, as a rule, on the lower parts of the stem both below and above ground, and are characterized by dark brown to black lesions or cankers, hence the origin of the German name "Schwarzbeinigkeit," or "blackleggedness." The first cankers are likely to develop below ground and not infrequently have their beginning at the lowermost part of the stem. Under favorable circumstances the disease works rapidly upward. On young succulent

stalks a lateral development of a lesion promptly results in a girdling of the stem, causing death. Under natural conditions the stem is seldom, if ever, first infected above ground. There is little tendency for the pathogen to migrate downward, even when stems are artificially inoculated at a point above ground. This fact is strongly emphasized by Smith's ('20) figures 193 and 199. The strong tendency on the part of the parasite to migrate upward along the main stem is shown in pl. 1, fig. 1. This photograph illustrates a rather striking, though not uncommon, case. Upon closer examination it will be found that the bacteria are largely confined to the parenchymatous tissues of the stem to the cortex and the pith. As it works upward in the pith this colorless parenchyma also becomes blackened and necrotic (pl. 1, fig. 4), easily observed by splitting the stem longitudinally. In this manifestation we have one of the most striking as well as reliable diagnostic features. The pith finally disintegrates, leaving the stalk more or less hollow. Frequently worms will be found feeding among the dead pith cells.

Lateral spread in large mature stems is very slight. Even though the stalk is severely diseased the plant may persist and mature a crop of tubers (pl. 2, fig. 3). Under favorable circumstances the infective agent moves through the stolons to invade the tubers. Sometimes the latter organs are infected at other points, invasion probably taking place for the most part through the lenticels, as has been experimentally demonstrated by Smith ('20). Diseased tubers frequently rot in the ground, the entrance of saprophytes bringing about the profound, slimy, putrid, soft-rot frequently observed. Death of the roots is secondary and follows a killing of the tops.

Tuber symptoms.—As implied above, stem-end infection of the tubers is most frequent under ordinary circumstances. As a rule, decay begins at the point of attachment of the stolon (pl. 2, figs. 3 and 4). Often, however, comparatively little decay is visible. Experiments carried out by the writer showed that the pathogen may be securely lodged in the tuber at the juncture with the rhizome, even though there are no signs of rot. He has also determined on many occasions that the blackleg parasite is the cause of more or less discoloration in the tissues associated with the vascular elements in the stem-end of the tubers. Such signs of the disease may readily be mistaken for symptoms of the *Fusarium* and *Verticillium* wilt diseases. In-

fection of the eye or bud end is less commonly found. A stem-end rotting of the Russet Burbank (Netted Gem) variety, due to *Bacillus atrosepticus*, is well known in many of the irrigated sections of the Northwest. It is most frequently seen at digging time, though it may be found earlier, especially if the latter part of the growing season is cool and rainy. Affected tubers are often found having pointed ends, a characteristic of some value, even though not proof-positive, since signs of the rot are more or less masked in freshly dug potatoes. Shrinkage of the affected tissues follows exposure of the tubers to drier surroundings and a brown to black discoloration shows through the skin. The presence of the rot is often exposed through breaking of the skin in the process of digging. Upon closer examination of affected tubers it will be found that the infection spreads more or less irregularly. In freshly dug tubers the rot may be described as a "soft rot" and is accompanied by a more or less putrid odor, but the blackleg as such is not to be confounded with this phase. Mixed infections, including the blackleg organism along with saprophytes and probably *Bacillus carotovorus*, are responsible for such conditions.

Dissection of tubers affected by the blackleg rot reveals the fact that the tissue may be involved to a considerable depth, extensive lesions often reaching the center. The color of necrosed tissues ranges from nearly normal to brown and black, but upon exposure to air these turn dark brown to black very rapidly (pl. 2, figs. 2 and 4). Advance of the rot is usually most rapid just beneath the epidermis (pl. 2, fig. 2), but in general spread in the tubers is not confined to any single region or tissue. Infected round tubers are often found to have a dark-colored hollow center.

A soft, cheesy rot caused by the blackleg bacillus is often seen among freshly dug tubers, though by some this type of rot is supposed to be due to "sunscald." The rot of tougher consistency found in affected tubers in transit or in storage is sometimes confounded with rots caused by *Fusarium* spp. Some very excellent illustrations of the tuber rot, including one colored plate, are those published by Shapovalov and Edson ('21).

Laboratory diagnosis.—To confirm the field observations it was necessary to make experiments in order to isolate the causal organism and prove its specific identity. Since sapro-

phytes quickly follow *Bacillus atrosepticus* in the tissues of both stem and tuber, it is often difficult to obtain the pathogen. By using extra precautions in carrying out the ordinary isolation methods one may obtain pure cultures of the causal organisms. A positive identification of the organism may then be made by careful comparison with the description of *B. atrosepticus* on p. 43.

ECONOMIC ASPECTS

Smith ('20) has come to regard the disease under consideration "as one of the most serious diseases of the potato." Earlier writers on the subject have unanimously emphasized its economic importance. Many cases are reported in the literature where upwards of 50 per cent of the plants in a field were destroyed by this disease. However, it appears that a loss of 5 per cent of the plants in a potato-growing district represents the usual amount of damage done by blackleg. In districts where the disease has been long known and where control measures have been intelligently applied, the losses have been greatly reduced, the average for the United States as a whole being about 0.5 per cent annually.

Dispersal and infection.—Probably the blackleg pathogen is most usually disseminated by the more or less general use of infected seed stocks. Morse ('16) mentions a striking illustration of this in a field in Idaho where he found blackleg which "undoubtedly came all the way from Scotland . . . in five years."

Without doubt the etiological agent is carried by wind and water, especially the latter, in the irrigated fields of the West. As pointed out by Smith ('20), invasion of the tubers through water-gorged lenticels can take place, but whether it does or not under natural conditions in the field is not known. The writer has planted healthy seed pieces in heavily infected soil but the disease did not develop in the plants thus grown. Similar results were recently attained by Shapovalov and Edson ('21). It appears to the writer that it still remains to be proven that the blackleg parasite can gain entrance to the vines or tubers through an unbroken epidermis.

The larvae of insects have been found by the writer working in and on the affected tissues, but there was no positive evidence that they were a tive agents in the dispersal of the disease. Von

Hegyí ('10) studied the disease on the continent of Europe, and since he found that every case of "blacklegged" stalks examined bore evidence of the attack of wireworms, he thinks insects are a positive factor in the invasion of the host. Paine ('17) believes that wireworms and biting insects are undoubtedly instrumental, under certain conditions, in introducing the parasite from the soil.

Geographical distribution.—The potato blackleg disease is now known to be widely distributed in the United States, Canada, and Europe. It was reported first in the northeastern United States and eastern Canada. Its spread, however, has been rapid, especially throughout the northern states. The disease has been reported from nearly every state in the Union. The records show, however, that it is essentially a cool-climate disease, therefore prevailing in more southerly districts only where potatoes are grown at high altitudes.

The writer finds that this disease in Montana and other sections of the Northwest is caused by the same organism known to prevail in the northeastern and North Central States. The recent work by Shapovalov and Edson ('21) proves that the same organism is widely distributed in the West.

Control.—Control measures should be inaugurated during the growing season. The writer strongly advocates the planting of a portion of the crop in a separate, if possible isolated, plot. This practice enables the grower to detect, remove, and destroy diseased hills at an early date.

Inspection and roguing of the plot (or field) should be begun soon after the plants are up and repeated at more or less frequent intervals throughout the growing season, especially during cool, rainy weather. Intelligent roguing will eliminate all hills where vines show any symptoms of the disease. Final inspections should be made just before and after digging. No tubers showing signs of rot or mechanical injury should be placed in storage. It is highly important that all seed stock be stored under favorable conditions. Slatted storage bins 8 or 9 feet high, 5 or 6 feet wide, and of any convenient length are ideal. In order to permit proper aeration of the potatoes, such bins should have false floors and must be separated from each other as well as from the walls of the cellar. The cellar should be fairly moist but provision should be made by proper regulation of ven-

tilators to prevent the atmosphere from becoming saturated. As nearly as practicable the temperature should be kept at about 38° F. in order to check sprouting. Butler ('19) has shown that potatoes kept at 40° F. will sprout after about 200 days, while at 35° F. sprouting is delayed indefinitely. Shapovarov and Edson ('19) showed that potato tubers which are in a wilted or softened condition when cut, due to the development of sprouts or improper storage, are very much more likely to be infected by fungous or bacterial parasites which exist in or may be introduced into the soil.

The seed pieces to be used for planting should be sorted and only sound, rot-free tubers used. Previous to cutting they should be treated in a 0.1 per cent corrosive sublimate (HgCl_2) solution or in a 1:240 formaldehyde dip. If the former is used, the tubers should be immersed for 1½ hours, then more of the dissolved mercuric chloride added at the rate of 1.5–2.0 grams (depending upon the amount of dirt accompanying the tubers) for each bushel treated. Avoid treating excessively dirty potatoes in the corrosive sublimate, and make up a fresh solution after having treated 50 bushels. If the formaldehyde solution is used the tubers should be immersed in the dip for about 1½ hours. Undoubtedly, a less lengthy immersion would be sufficient to kill all contaminating blackleg germs present. When the pathogen is lodged internally treatments sufficient to kill it would probably kill the buds also.

When and where practicable, planting should be done late enough to avoid having the crop sprout and struggle along in a cold, damp soil, since it has been shown that these conditions facilitate the development of the blackleg parasite. While it has not been conclusively proved that the blackleg parasite does not overwinter in the soil at times, it is nevertheless advisable to practice rotation of the crop, since some of the worst enemies of the potato accumulate in the soil and persist therein for long periods of time.

Tests made by the writer show that the blackleg bacillus is resistant to considerable extremes of cold. Test-tubes containing about 15 gms. of soil were thoroughly autoclaved and the sterility of the soil carefully tested before inoculation. Finally, a series of cultures thus prepared were inoculated with a freshly invigorated strain of the organism and placed out of doors for 24

hours. During this time the official minimum was -28° C. and the maximum -6.7° C. Soil cultures similarly inoculated were incubated at 28° C. The cultures exposed out of doors were placed on ice upon being brought in, in order to prevent excessively rapid thawing. Finally transfers were made to nutrient broth, and agar slants from all the cultures included in the experiment. The organism was recovered from all the cultures. The subcultures made from soil which had been frozen for 24 hours appeared to be as vigorous as any.

There are few satisfactory data on varietal resistance of potatoes to the blackleg pathogen. An early contribution to this phase of the problem was made by Appel ('03), who concluded that the thick-skinned, late varieties which were being grown in Germany were more resistant to the disease than the thin-skinned, starch-poor, early varieties. Morse ('17) reported that the Irish Cobbler and Green Mountain varieties were particularly susceptible to the disease. The Early Ohio, an early variety, is notably susceptible. The Russet Burbank and the Idaho Rural, varieties widely grown in the West, are commonly found affected with blackleg.

THE ETIOLOGICAL AGENT

The writer's studies on the etiology of the potato blackleg disease were begun in Montana some years ago. By the close of the summer of 1915 extended observation led to the assumption that the blackleg disease in Montana was quite similar to that occurring in some of the eastern states. In the meantime some 30 isolations were made at the Montana Experiment Station from potatoes affected with blackleg. The pathogenicity of the strains thus isolated was thoroughly tested for the second time in 1916, and a majority was found to be pathogenic and capable of causing typical symptoms of the disease. Comparative studies of the morphological, cultural, and physiological features of some of the above-mentioned strains were made by the writer. The results obtained led to the conclusion that all were essentially alike. An extension of these preliminary investigations was made, and cultures of the blackleg bacillus from Maine, Minnesota, and one from eastern Canada received under the name *B. solanisaprus* Harrison, were cultivated and compared with one another and with observations made previously on the Montana strains. While the observations made at this time were

not extended, the writer was led to assume that a single organism caused the potato blackleg disease in North America.

A study of the literature was begun, and it was soon learned that at least 4 different species of *Bacillus* had been described by as many authors as the cause of the disease under consideration.

By comparing the following descriptions it will be seen that the "species" in question do not differ markedly. In the summaries presented no attempt is made to follow the original form, but accuracy of statement is preserved.

Bacillus atrosepticus van Hall.—The following was prepared from a translation¹ of van Hall's ('02) original description. It bears comparison with that prepared for Morse ('17) by Dr. R. de Zeeuw.

Morphology: *B. atrosepticus* is a rod-shaped bacillus, occurring for the most part singly, rarely in pairs, in 2-day-old bouillon cultures at 27°C. Size variable; length 0.8–1.6 μ , breadth 0.2–4 μ . Many zoogloea of 4–10 organisms. The bacteria are very active in a 24-hour culture of distilled water plus 0.025 per cent potassium phosphate and 0.25 per cent asparagine (27°C.). Material stained by Loeffler's method. Length of flagella 10–15 μ . Gram negative.

Cultural characteristics: Gelatin liquefied, rapidity variable. Growth on malt gelatin and malt agar very weak. Milk coagulated. Growth best at top of meat agar stab.

Physiology: Growth very strong at 27°C. Thermal death point 51–52°C. Facultative anaerobe. Reduction of methylene blue weak. Nitrates reduced to nitrites. Sodium selenite reduced rapidly. No diastatic action. No indol production. No H_2S produced in broth cultures. The organism is a weak gas producer (except when mannite is present). Gas produced from glucose, saccharose, and mannite in a medium made by adding to "due water" (a filtered water) 0.025 per cent K_2HPO_4 , 1 per cent peptone and 3 per cent of sugar. No gas from lactose and glycerin in same medium. Growth slow at first in bouillon acidified to a reaction of 0.5 per cent normal with citric and malic acid. Organism grows poorly when transferred from dried cultures: thought not to be resistant to drying. Pathogenic to potato stems and tubers. Index No. 5312-32120-2121.²

Bacillus phytophthorus Appel.—The following is summarized from Appel's ('03) diagnosis:

¹My thanks are due to Dr. J. C. Th. Uphof for assistance rendered in translating Part III of van Hall's ('02) dissertation.

²"Index Number" digits throughout this paper are arranged according to the Descriptive Chart indorsed by the Society of American Bacteriologists, Dec. 30, 1920.

Morphology: *Bacillus phytophthorus* a fairly thick, colorless, rod-shaped organism. Breadth averages $0.8\ \mu$, length $1.2\text{--}1.5\ \mu$. Usually occurring singly, occasionally in pairs. Motile by means of (usually six) peritrichic flagella, stained by Peppler method. Gram negative.

Cultural features: Liquefaction of gelatin saccate. Gelatin colonies small, yellow-white, with entire margin. Agar colonies small, glistening, bluish-opalescent. Growth on agar stroke rapid, moderate to abundant. Upon sterilized potato plugs a weak honey-yellow growth develops. On sterile raw potato slices, growth and browning in 15-18 hours after inoculation. In 2 days middle line sinks and surrounding tissue becomes dark on upper surface. A liquid is pressed out. Growth in potato juice rapid and abundant. Growth in nutrient broth moderate, turbid and sediment.

Physiology: Gas from sucrose and maltose. No gas in shake cultures of sugar agar. Nitrates reduced to nitrites. Milk coagulated slowly, acid (?) curd. Pathogenic to potato stems and tubers. Index No. 5112-32120-1111.

Smith's description of *B. phytophthorus* Appel.—Inasmuch as Appel's ('03) original diagnosis of his species is meager, the following compilation of E. F. Smith's ('20) description is presented:

A white, rapid-growing, non-sporiferous, Gram-negative, motile, peritrichiate-flagellate, promptly liquefying, nitrate-reducing, aerobic and facultative anaerobic, acid-forming, gas-forming, milk-curdling (by formation of an acid), dry-air-sensitive, rod-shaped or filamentous schizomycete, forming quickly on agar plates circular, grayish white, well-developed colonies; on very thin-sown gelatin plates characteristic, rapid-growing, big, circular, opaque white colonies. Organism white on most media but on Soyka's milk rice it is pale pinkish cinnamon. Bouillon clouded very quickly and gelatin stabs develop a prompt funnel of liquefaction. The organism does not form indol, and does not grow in Cohn's solution. It produces a non-volatile acid from dextrose, saccharose, lactose, galactose, and maltose, and small quantities of gas from inosit (muscle sugar), lactose, and mannite. Pathogenic to potato shoots. Index No. 5312-32120-1212.

***Bacillus solanisaprus* Harrison.**—Summary given herewith is compiled from Harrison's ('07) original description:

Morphology: Bacillus of variable size. From 24-hour-old beef agar spores not seen; flagella peritrichic; organism Gram negative.

Cultural features: Growth in agar abundant, glistening, opalescent; persistent growth on potatoes; uniform growth in gelatin cultures, filiform liquefaction; strong growth in nutrient broth; ring in surface growth; clouding strong, fluid turbid, fine sediment; growth in gelatin colonies slow, round to elliptical; agar colonies round to lenticular, shiny, entire.

Physiology: Prompt coagulation of milk, extrusion of whey in 3 days, a few gas bubbles visible; litmus milk acid. Strong growth in Uchinsky's solution. In fermentation tubes gas was produced in bouillon containing mannite and lactose; growth in closed arm, with production of acid in each of the following: dextrose, saccharose, lactose, maltose, glycerin, mannite, and levulose. Nitrates in nitrate broth were reduced, nitrites present, indol production moderate to feeble; vitality on culture media long; thermal death point 54° C., optimum temperature for growth $25-28^{\circ}$ C., growth slight at 37° C.; maximum temperature for growth 37.5° C., minimum temperature for growth about 0° C. Pathogenicity proven to following vegetables: potato, tomato, Jerusalem artichoke, and others; also to living plants of potato, tomato, common red pepper, and slightly in cucumber and physalis. Index No. 5312-32120-1212.

Bacillus melanogenes Pethybridge and Murphy.—The account presented below is summarized from Pethybridge and Murphy's ('11) original diagnosis:

Morphology: Vegetative cells $0.7-0.9 \mu \times 1.3-1.8 \mu$, found most frequently in pairs. Flagella peritrichic (fewer than in *B. solanisaprus* Harrison). No endospores. Gram's stain negative.

Cultural features: Nutrient broth clouded, more or less sediment, fluid turbid, no pellicle. Organism did not form a distinct ring on surface of potato juice. Gelatin colonies round. Growth best at top in gelatin tubes, some growth along needle track. Gelatin liquefied. On potato plugs growth abundant, and chromogenesis yellowish.

Physiology: Facultative anaerobe. Milk acid curd, curd not very compact. No indol produced, nitrates reduced to nitrites with formation of gas. Diastatic action on starch. Acid and gas in fermentation tubes, in broth plus glucose, lactose, and saccharose. No acid from glycerin. Pathogenic to potato.

They comment upon the marked resemblance of their organism to *B. phytophthorus* Appel and state that they were tempted to regard it as a variety of the latter. Index No. 5312-3211?—1111.

After comparing the contributions of these earlier investigators, and in the light of the observations which I had previously made, I planned a thoroughgoing study of the relationship of the blackleg pathogen. To this end it was thought best to make careful comparative studies of a number of strains, including if possible the 4 species described above.

Source and history of the cultures used.—Some difficulty was experienced in obtaining authentic and viable subcultures of the organisms described by van Hall, Appel, Harrison, and Pethybridge and Murphy. Cultures of these organisms, however, were

finally obtained.¹ These were supplemented by cultures of the blackleg parasite isolated in Montana, Minnesota, and Maine. For the sake of convenience all of the subcultures collected were designated by a number. In order to facilitate matters, all records of studies made on the strains selected were kept under an assigned number.

Below is given a brief note on the source and history of the several cultures used more or less extensively throughout the writer's comparative studies. The first 5 enumerated were from isolations made in the Botany Department at the Montana Experiment Station and were selected from isolations made from potatoes affected with the blackleg disease obtained in some of the more important potato-growing sections of that state.

No. 160.1. Isolated 8/14/15, from tuber affected with black rot. Material from near Bozeman, Mont. Pathogenicity established. Still virulent late in 1917.

No. 170.3. Isolated 8/22/15, from tuber affected with black soft rot. Material from near Lewistown, Mont. Pathogenicity established. Still virulent late in 1917.

No. 180.2. Isolated 7/24/16, from black, cortical lesions on stem. Material from near Kalispell, Mont. Pathogenicity established. Still virulent late in 1917.

No. 183.2. Isolated 8/27/16, from black cortical lesions on stem. Tubers on same vine affected with blackleg rot. Pathogenicity established. Still virulent late in 1917.

No. 187B.1. Isolated 9/2/16, from tuber with superficial black rot lesions. Pathogenicity established. Still virulent late in 1917.

No. 191. Kindness of W. J. Morse. His "*B. phytophthorus* from Appel." Pathogenicity not established by the writer. Morse found it non-pathogenic.

No. 193. Kindness of W. J. Morse. His "III A," isolated by him in Maine, Aug., 1908. Found by the writer to be pathogenic. Still virulent late in 1917. Used by Morse in his studies ('17).

No. 194. Kindness of W. J. Morse. His "SE." Isolated by him in Maine, Aug., 1908, from a "potato stem showing a very rapid soft rot." Found by the writer to be pathogenic. Still virulent in late 1917. Used by Morse ('17).

¹ My thanks are especially due Dr. W. J. Morse and Dr. B. M. Duggar for very material assistance rendered in supplying certain of the cultures used in this work.

No. 195. Kindness of W. J. Morse. His "II P." Isolated by him in Maine, Aug., 1908, "from typical blackleg plants". Found by the writer to be pathogenic to potato tubers. Still virulent late in 1917. Used by Morse ('17).

No. 196. *B. solanisaprus* Harrison. Kindness of W. J. Morse, who used this strain in his studies ('17). The culture was originally procured from S. F. Edwards, Ontario Agr. Coll., in March, 1909. Found by the writer to be pathogenic to potatoes. Virulence in 1917 good.

No. 197. *B. atrosepticus* van Hall. Kindness of W. J. Morse. Previously studied by him ('17). This strain Morse "received under that name from Kral's laboratory¹ in 1910. . . . At first it showed weak pathogenicity to potato tubers, and repeated inoculations to growing stems failed to produce the disease until the present summer [1916]." Found by the writer to be pathogenic, though, if anything, less strongly so than some of the others. Still virulent late in 1917.

No. 198. *B. melanogenes* Pethy. and Murphy. Kindness of W. J. Morse, who "received this from Dr. Pethybridge himself in 1911. . . . It also has produced active decay of tubers and blackleg of the stem upon inoculation." Used by Morse ('17). I found it to be pathogenic. Still virulent in Nov., 1917.

No. 200. "*B. phytophthorus*" from Minnesota. Kindness of E. C. Stakman. Isolated 1915. The writer found it to be pathogenic to potato tubers. Still virulent in 1917.

No. 201. *B. phytophthorus* Appel. Kindness of B. M. Duggar who obtained the culture from E. F. Smith. The latter procured the strain from Dr. Aderhold in Berlin about 1906. It is presumed that Smith's culture was a transfer from Appel's original culture. The writer found it to be pathogenic. Still virulent in 1917.

No. 202. "*B. solanisaprus*." Kindness of D. H. Jones who wrote that "the transfer purports to be from original strain of *B. solanisaprus* . . . received in Oct., 1916, from the American Museum, to which Dr. Harrison had formerly sent a culture." Found by the writer to be pathogenic. Still virulent in late 1917.

II. COMPARATIVE STUDIES OF CAUSAL ORGANISMS

Invigoration of cultures.—The cultures used by the writer were growing on beef agar at the time the comparative studies presented below were begun. Some had been previously cultivated on agar and some in nutrient broth for greater or less length of time. To begin with, all were plated out to insure purity of the culture and then each one was invigorated by trans-

¹ Kral's Bakteriologisches Laboratorium, Prague, Austria.

ferring to nutrient broth and cultivating at 25–27° C., on 3 consecutive days. All the cultures so treated responded promptly.

Pathogenicity.—Testing the pathogenicity of the strains to be used proved to be a considerable task. A number of trials were made before satisfactory proof was obtained of the pathogenic character of all of the strains in use. In the first series of tests potato tubers were used. On the whole, sound, smooth tubers of some thin-skinned variety, such as Early Ohio, provided the most satisfactory material for artificial inoculation experiments. Thick-skinned, late varieties, as the Russet Burbank, can be used. Tubers with a relatively high sugar content proved to be the best for this work (cf. Carbohydrate Utilization, page 45). On this account it is well if possible to obtain growing tubers, or else those from storage which are held till sprouts begin to develop.

Method.—Selected tubers were thoroughly washed by scrubbing in warm water, then soaked 3–4 hours in clean warm tap water, finally disinfected by immersing for 15 minutes in a 0.1 per cent aqueous solution of mercuric chloride. Immersion in this disinfectant for a longer period of time is not necessary. Moreover, it was found when the tubers were soaked for 1½–2 hours that the chemical remained in the outer layers of the skin, to the extent of making it difficult, if not impossible, to remove all of it by washing. Not infrequently the effects of a long soaking were manifest by inconspicuous lesions in the surface layers. Clean ground-glass plates, watch-glasses, and bell jars were made ready and sterilized in advance by washing in the bichloride solution. The tubers were then removed from the disinfecting solution, thoroughly washed in sterile tap water, and placed on watch glasses under the bell jars.

Preliminary experiments demonstrated the desirability of maintaining a nearly saturated atmosphere surrounding the tubers during incubation for 4 or 5 days. This requirement was easily satisfied by lining the inside wall of each bell jar with sterilized filter-paper wet with sterile water. If the atmosphere of the room is dry and cool it will be found best to place the inoculation chambers in an incubator kept at 22–25° C. All but one of the tubers under each bell jar were inoculated. Inoculations were made in a clean damp culture room. It was accomplished by placing a drop of the suspension from a recently in-

vigorated culture in a depression, usually near an eye, and stabbing through the inoculum 3 or 4 times with a sterile needle. The control tubers were pricked in similar places with a sterile needle. Inoculation courts were marked with an indelible pencil by tracing a good-sized circle around the spot where the tuber was pricked. Check courts were marked by a square.

Observations and conclusions.—The inoculation sets thus prepared were closely observed every day for 10 days, then with less frequency for over a month.

The number of days elapsing before signs of infection were visible varied, in the several experiments performed, from 2 to 6 days. Infection was in general more prompt where tubers having a high sugar content were inoculated, where the set incubated at around 23° C., and where the moisture content of the atmosphere surrounding the tubers was kept high. Observations made by the writer on the pathogenicity of several strains of the potato blackleg parasite suggest the conclusion that the "incubation period" in plants is markedly influenced by environmental conditions. The "incubation period" did not appear to be marked by a definite "turning point," as is the case in many animal diseases.

The writer concluded that all the strains being tested were pathogenic (cf. tables I and II). The nature and development

TABLE I
TABULAR SUMMARY
SOURCE AND HISTORY OF CULTURES USED

Cult. No.	Isolated	Source	Pathogenicity		Comparative virulence
			Stems	Tubers	
160.1	Tuber 8-14-'15	Montana	Marked	Strong	Average
170.3	Tuber 8-22-'15	Montana	Marked	Strong	Average
180.2	Stem 7-24-'16	Montana	Strong	Marked	Average
183.2	Stem 8-27-'16	Montana	Marked	Marked	Average
187B.1	Tuber 9-2-'16	Montana	Strong	Strong	Strong
193.	"Plant" Aug., '08.	Maine, Morse's IIIA	Strong	Strong	Average
194.	"Stem" Aug., '08.	Maine, Morse's SE	Yes	Yes	Average
195.	"Plant" Aug., '08.	Maine, Morse's IIP	Strong	Yes	Average
196.	Before Mar., '09	<i>B. sol.</i> Ontario, Can.	Marked	Marked	Average
197.	Before 1910	<i>B. atro.</i> Kral's lab.	Marked	Yes	Average
198.	By P. & M. 1911.	<i>B. mel. fr.</i> Pethy. Ireland	Yes	Yes	Low
200.	"Blackleg" 1915	Minnesota	No data	Yes	Low
201.	Before 1906	<i>B. phyto. fr.</i> Germany	Marked	Marked	Strong
202.	About 1907	<i>B. sol.</i> Am. Museum	No data	Strong	Average

of rot lesions in the tubers varied considerably under the different conditions of the artificial inoculation experiments. Lateral spread from the points of inoculation seldom took place to the extent it does in tubers invaded at the stem end and under natural conditions. The depth to which the lesions extended was usually to a point considerably beyond the depth of the needle prick, occasionally to the opposite side of the tuber. The character of the rot developed in the tubers inoculated artificially varied somewhat. Sometimes the affected tissues were quite moist and soft. Again, the tissues attacked were quite dry, being more or less spongy or cheesy in texture. Diseased material exhibited a more or less putrescent odor but in no case did inoculation with pure cultures effect a profound, gray, slimy, obnoxious, gaseous, soft rot. Variations, such as those mentioned, appeared to be associated to a considerable degree with (1) the available sugar content of the tuber, (2) the moisture content of the tuber, and (3) with certain environmental factors, notably temperature and humidity. The lesions in immature potatoes were generally more profound and the affected tissues softer than those in ripe tubers, particularly of a late, hard, starch-abundant variety. A drop of inky black liquid was exuded at the points of inoculation in many cases. Occasionally a slight bulging of the tissues occurred at these points. When affected tubers were cut or broken open the diseased tissues were characteristically dark-colored, becoming brown or black upon exposure to the air.

Additional observations.—To extend the above observations on pathogenicity of the various strains at hand, the artificially inoculated tubers were cut into seed pieces and planted in pots in the greenhouse. In many instances the seed piece was entirely consumed by the blackleg rot before sprouts started. In a few cases, however, sprouts developed, only to be rapidly invaded by the pathogen and killed. The case of culture 160.1 is typical. Four stalks emerged from the ground but by the time they had grown to be from 3 to 6 inches tall (30 days after planting the seed pieces), the parasite invaded the stem tissues. The infection extended rapidly upward. At first the diseased tissues (cortical) appeared water-soaked. The tissues became dark-colored over night, and in a day or two the plants were prostrated (pl. 1, fig. 3). What proved to be a pure culture (plates) was recovered from this material. This was later used for inoculation work and found to be pathogenic in tubers.

Some experiments on the pathogenicity of the potato black-leg parasite performed at later dates furnish further proof of the pathogenicity of the strains under observation, and will be reported here. A brief survey of the results obtained in one of these experiments (II of 7/21/17) appears in tabular form below (table II). Tubers of the Russet Burbank variety were used. They had been kept in storage nearly a year. The tubers were washed and disinfected in the manner described previously. A number of selected tubers were then cut into seed pieces, each weighing 2 to 3 ounces. Three seed pieces were then inoculated with each of the several strains of the organism at hand. Two were planted in pots, the third being placed in a moist chamber kept at room temperature (20–24° C.)

TABLE II
PATHOGENICITY OF STRAINS OF *B. ATROSEPTICUS* EMPLOYED

Strain No.	Pathogenicity to seed pieces planted in soil	Pathogenicity to seed pieces kept in moist chamber
160.1	Positive (sprouts involved)	Negative (no infection, piece dry)
170.3	Positive (no sprouts)	Negative (no infection, piece dry)
180.2	Positive (no sprouts)	Positive (black rot)
183.2	Positive (no sprouts)	Positive (black rot)
187B.1	Positive (no sprouts)	Negative (no infection)
191.	Negative (sprouts)	Negative (non-pathogenic)
193.	Positive (sprouts)	Positive (black rot)
194.	Negative (sprouts)	Positive (black rot)
195.	Negative (sprouts)	Negative (no infection)
196.	Positive (no sprouts)	Positive (black rot)
197.	Negative (one sprout)	Positive (black rot)
201.	Positive (no sprouts)	Negative (no infection)

Evidently all the strains used were still virulent.

Among other things this experiment shows that the black-leg organism may destroy infected seed pieces before sprouts develop. In other cases the infectious material may be rendered innocuous or even killed by drying and by other agents. These conclusions are emphasized by experiments performed and reported above.

An experiment begun in August, 1917, showed that the strains previously listed were pathogenic to potato tops; that is, all introduced artificially gave rise to typical blackleg lesions in the stems and in leaf petioles.

A series of experiments was planned during the summer of 1917 in order to throw light on the question of whether potatoes planted in a soil contaminated with the blackleg parasite would

become affected with the disease. Only one trial was made, and while the results obtained do not in themselves warrant drawing a conclusion, they are of interest when compared with those reported by Shapovalov and Edson ('21). Ordinary seed pieces from sound, hard, healthy potatoes were planted in soil which had been previously heavily seeded with the blackleg germ by planting with tubers affected with the disease. The seed pieces sprouted and the plants therefrom grew to be of good size. No signs of infection by the blackleg germ were to be found in the seed piece, on the roots, or in the tops.

METHODS AND MEDIA

Certain of the bacteriological methods recommended by the American Public Health Association ('12) were followed as closely as possible by the writer in the prosecution of his studies of the cultural characteristics and physiology of the blackleg strains selected for comparison. In some cases this was not practicable or even possible on account of the fact that certain reagents were not available, due to conditions brought on by the World War. For this reason, a rather complete statement is given below of the media and methods employed, thus making it possible to duplicate both.

The present-day methods for determining H-ion concentration of bacteriological media were not in general use at the time the writer carried out his comparative studies. The ordinary media used were titrated with N/20 NaOH, using phenolphthalein as an indicator. In most cases the reaction of the medium was not adjusted. Wherever stated the reaction is given as "+7" etc., where the addition of 7 cc. of normal alkali per 1000 cc. would render the medium neutral to phenolphthalein. A "O" indicated that the medium as used was neutral to phenolphthalein. The precaution of having all flasks, beakers, test-tubes, etc. thoroughly clean was taken at all times.

Media.—Steps in the preparation of the media are briefly indicated below:

NUTRIENT BROTH

(1) To 1000 cc. distilled water add 50 gms. "Bacto" Beef Extract; (2) heat the mixture gradually up to 70°C. during about 1 hour; (3) boil for a few minutes to coagulate precipitable proteins; (4) make up the water lost by evaporation; (5) filter through layers of clean cheese-cloth; (6) cool to about 60°C. and add 1 per cent "Difco" peptone, $\frac{1}{2}$ per cent NaCl, and dissolve; (7) filter through

paper; (8) titrate, using phenolphthalein as an indicator; (9) determine reaction and adjust if necessary; (10) sterilize in autoclave at 15 pounds for 15 minutes.

BEEF AGAR

(1) Nutrient broth (steps 1-6) is brought to a boil; (2) while boiling add 1 per cent "Bacto" agar, sifting it over the surface; (3) when agar is completely dissolved, cool and clarify in usual manner; (4) determine reaction and adjust if necessary; (5) filter; (6) sterilize in autoclave.

GELATIN

(1) To nutrient broth already prepared (steps 1-6), add 10 per cent "Bacto" gelatin and dissolve at a low temperature (60-70°C.); (2) determine reaction and adjust if necessary; (3) filter, tube, and sterilize at 15 pounds for 15 minutes.

DUNHAM'S SOLUTION

Make a paste of 10 gms. peptone and 5 gms. salt in a small quantity of water; then add sufficient water to make total used 1000 cc. Heat in flowing steam for $\frac{1}{2}$ hour, then boil for 10 minutes over free flame. Make up water lost by evaporation. Filter through paper and tube. Sterilize in autoclave at 15 pounds for 15 minutes.

CARBOHYDRATE BROTH

(1) To nutrient broth already prepared (steps 1-6) add 1 per cent by weight of the chemically pure carbohydrate; (2) determine and record reaction; (3) filter, tube, and sterilize in autoclave at 15 pounds for 15 minutes.

MILK

The milk used was freshly drawn into a sterile bottle. After standing about an hour it was centrifuged and the cream skimmed off. Samples were titrated, using phenolphthalein as an indicator, and the reaction was determined to be "+20". The reaction was not adjusted. The medium was tubed promptly and sterilized by exposing tubes in flowing steam for 20 minutes on 3 consecutive days.

LITMUS MILK

To part of the fresh milk (see above), was added 2 per cent by volume of a saturated aqueous litmus solution (Merck's reagent, 1 gm. to 15 cc. H_2O). This medium was tubed immediately and sterilized by exposure to flowing steam for 20 minutes on 3 consecutive days. Both milk media were kept 4 or 5 days before being inoculated.

RAW POTATO PLUGS

A considerable number of raw potato plugs were prepared, observing all possible antiseptic precautions. The lot was kept several days in order to insure the removal of any contaminations that might appear.

RAW POTATO DISKS

Sterile disk-shaped slabs from raw potatoes were successfully prepared in the same manner as the preceding lot. They were placed in sterile Petri dishes on a piece of sterile wet filter-paper. On the whole, these disks proved to be more usable than the raw potato plugs.

COHN'S SOLUTION

Dissolve 5.0 gms. KH_2PO_4 , 5.0 gms. MgSO_4 , 10.0 gms. $(\text{NH}_4)_2\text{C}_4\text{H}_4\text{O}_6$, and .5 gm. $(\text{Ca}_3(\text{PO}_4)_2)$ in 1000 cc. distilled water, and filter through paper. Sterilize in autoclave at 15 pounds for 15 minutes.

USCHINSKY'S SOLUTION (MODIFIED)

The modified solution was made up according to the formula given by Smith ('05).

EHRlich's INDOL TEST SOLUTION

Solution I.—Para-dimethyl-amido-benzaldehyde, 4.0 gms.; 95 per cent alcohol, 380 cc.; HCl (conc.), 80.0 cc.

Solution II.—Saturated aqueous solution of potassium persulphate.

To about 10 cc. of the liquid culture (preferably in Dunham's solution and 10 days old), add 5 cc. of Solution I, then 5 cc. of Solution II, using separate clean pipettes. Shake the mixture. The reaction may be accelerated by heating to about 70°C . The appearance of a red coloration which increases in intensity indicates the presence of indol.

NITRATE BROTH

Filter, tube, and sterilize in the autoclave 1 gm. peptone ("Bacto"), 0.2 gm. potassium nitrate cp. (Merck), and 1000 cc. distilled water.

REAGENTS AND TEST FOR NITRITES

Test for nitrites in nitrate broth culture on fifth day by adding: (1) one cc. of a 1 per cent potato starch water; (2) one cc. of a freshly prepared KI water (i.e., KI, 0.2 gm. in 50 cc. distilled water); (3) a few drops of strong H_2SO_4 , (i.e., conc. H_2SO_4 , 1 part, distilled H_2O , 2 parts). The appearance of a blue-black coloration in 10 to 15 minutes was taken as positive evidence of the presence of nitrites.

DETAIL OF COMPARATIVE STUDIES
MORPHOLOGY

A detailed study of the morphological characteristics of the several strains at hand was not attempted by the writer. However, it was determined that each was a small, short, actively motile, rod-shaped organism, easily stained by the ordinary aniline dyes. Pairs of the organism occurred in preparations made from fresh cultures on agar slants. No spores were found in any of the cultures, and all were determined to be Gram negative.

Cultural features and physiology.—In prosecuting the general plan outlined for making comparative studies of the cultural features and physiology, it was deemed wise to give particular attention and painstaking effort to the study of those items which are most commonly made use of in establishing bacterial species.

In recent times the response of the microorganism, as well as the reactions set up by it, in the presence of carbohydrates, has been stressed in attempts to distinguish between closely related species. On this account, and because of the fact that earlier students of the potato blackleg parasite report a great variety of conclusions as to the ability of this organism to ferment certain carbohydrates, the writer made exhaustive studies upon the gas- and acid-producing function of the blackleg bacillus. Details of the comparative studies appear on the several tables presented herewith and in the paragraphs below.

Nutrient broth:— Clouding prompt, persistent; medium becoming slightly turbid; sediment compact, granular, somewhat viscid; surface growth a ring, though under certain circumstances a light pellicle may form; no chromogenesis; odor absent; color of medium unchanged.

Earlier students of the blackleg bacillus did not wholly agree as to cultural features exhibited by this organism when grown in nutrient broth. The several strains studied were grown in 3 nutrient broths: (A) "Bacto" beef and "Bacto" peptone, (B) Liebig's beef extract and Witte's peptone, (C) fresh meat infusion and Witte's peptone, and all incubated at 27–28° C. All (except No. 191) were very similar in their response in nutrient broth. In those grown in "Bacto" beef and "Bacto" peptone (except Nos. 191 and 198) the surface growth was a ring, clouding was moderate, sediment granular-viscid, the medium slightly turbid, and no color. In the case of 191 the surface growth was a membrane, and in 198 a light pellicle with moderately strong clouding. When the strains were grown in Liebig's beef extract and Witte's peptone the surface growth was a ring (except No. 191 which was a membrane), clouding moderate, sediment granular viscid, medium slightly turbid, chromogenesis none. In those grown in fresh meat infusion and "Bacto" peptone, except Nos. 160.1 170.3, 187B.1, 191, 197, and 201, the surface growth was a pellicle, the clouding strong, the sediment granular-viscid, medium slightly turbid, and chromogenesis none. Nos. 160.1 and 187B.1 showed the surface growth a wide ring, and

moderate clouding; No 170.3 and 197, moderately strong clouding; No. 191, the surface growth a membrane with moderate clouding; and in No. 201 the surface growth was a light pellicle.

Observations recorded upon the cultural features exhibited by the different strains (except No. 191) as grown in these broths suggest a clue as to the disagreement between all who have cultivated the blackleg bacillus as to the character of the surface growth. It was found that the clouding was stronger and the surface growth heavier (a pellicle) in the nutrient broth made from a fresh meat infusion and "Bacto" peptone than in either of the other two nutrient broths used. Moreover, the growth response in the broth made with "Bacto" beef and "Bacto" peptone was somewhat more luxuriant than that in the beef extract and Witte peptone broth. The writer concludes that strain No. 191 is the only one of the 12 studied in broth that exhibits dissimilarity.

Agar stroke.—Growth moderate, filiform, somewhat raised, with smooth surface and glistening luster, translucent and somewhat iridescent, non-chromogenic; consistency butyrous, some quite viscid; odor absent; medium (color) unchanged.

On agar slants the cultural features were identical for the strains studied, except No. 191, where the form was spreading and effuse, the surface smooth, and luster dull.

Potato (cooked).—Growth moderate to abundant, filiform at first, spreading after a few days, slightly raised and flat, glistening, with a smooth to slightly contoured surface; chromogenesis yellowish white; odor not strong nor characteristic; color of medium slightly browned.

Special attention was devoted to the cultivation and study of the cultural features of the blackleg strains on sterilized potato plugs. This was done because of the importance attached by some authors to the cultural characteristics on this medium, in the differentiation of their "species" from others. All the strains studied comparatively by me were quite alike (except No. 191), in so far as can be detected from a study of the growth and cultural features on sterilized potato plugs. In No. 191 the growth was scanty, form spreading, surface smooth, dull, and colorless.

Potato (raw).—Raw potato slabs aseptically prepared as outlined in a preceding section were used. Preliminary tests indicated that infection and growth by the organism took place when the atmosphere of the culture chamber (Petri dish) was moist and when incubated at about 26° C.

The following strains were tested: Nos. 160.1, 170.3, 180.2, 183.2, 187B.1, 191, 195, 196, 197, 198. All responded and grew very similarly, except No. 191. In the case of the other 9 strains a considerable browning of the tissue approximate to the lines of inoculation was plainly visible at the end of the first day. A small quantity of dark brown to black liquid collected adjacent to the line of growth. By the end of a week abundant development of the cultures had taken place, forming a grayish white slime surrounded by a dark-colored border. The tissues at the margin of the growth were dark brown to black in color. Underneath the bacterial growth they were somewhat collapsed, so that the culture was growing in a shallow depression. No growth took place on the uninoculated slabs kept as controls.

Agar colonies (tubes inoculated at 28° C.).—Colonies rather small (3–15 mm. in 15 days); growth moderately rapid; form round to somewhat irregular; surface smooth; elevation raised and flat; edge entire, becoming undulate or lobed; internal structure, when magnified about 50 times, finely to coarsely granular; chromogenesis none; submerged colonies small, biconvex to ovoid, or nearly round; color of medium unchanged.

In strain No. 191, the colonies were large; surface smooth; elevation effuse; edge erose; and internal structure fine. This was the only one of the 11 planted in agar plates that developed markedly different features. In fact, one is led to the belief that no more than 1 species could possibly be represented by the 10 strains in question.

Gelatin colonies (incubated at 18–23° C.).—Form round to somewhat irregular; edge entire, often becoming undulate; liquefaction first noticeable, as a rule, about 24 hours after planting, proceeds rapidly, and is saucer-shaped; chromogenesis none.

Here again the cultural features and the physiological response as manifested by liquefaction were identical for all strains, except No. 191. In this case there was no liquefaction, and the colonies were larger. At about 21° C. colonies developed quite promptly in gelatin, and averaged 5–20 mm. in size, which was considerably larger than the agar colonies.

Gelatin stab.—Growth best at top; line of puncture filiform, often slightly beaded; liquefaction begins in about 24 hours and at 20° C., at first crateriform or napiform, becoming infundibuliform or strati-form, complete as a rule at end of third week, when culture is aerated by shaking occasionally; the liquid gelatin becomes cloudy and often a ring; sometimes a pellicle develops on the surface.

With a single exception (No. 191) the strains cultivated by stabbing gelatin plugs grew and responded almost identically as may be seen by glancing at table III.

Milk.—Fresh milk (+20) incubated at 28°C. is promptly coagulated. The amount of acid developed by the organism increases steadily for 18 days at least. Peptonization slow, begins in 2–3 days. In similar cultures to which litmus was added, a marked bleaching of the indicator took place by the end of the eighth day, as a rule. The addition of 2 per cent (by volume) of a saturated solution of Merck's purified litmus had no noticeable inhibitory effect upon the growth of the strains of the blackleg bacillus tested.

It may be noted that considerable difference appears in the descriptions published by earlier authors as to the cultural features and physiology of the blackleg bacillus when grown in milk. The writer failed to discover any real differences among the strains cultivated by him except in the case of strain No. 191 which, as has appeared heretofore, is non-pathogenic and otherwise quite different in its cultural features and physiology (see table III).

Uschinsky's solution.—Growth copious; fluid not viscid.

In this solution in which the nitrogen is supplied by organic compounds (see p. 28) all strains cultivated developed similarly except No. 191, which grew more copiously than the others with the development of a pellicle upon the surface of the medium.

TABLE III
CULTURAL FEATURES AND PHYSIOLOGY

GELATIN STAB*		REACTION 0	INCUBATED AT 20° C		
Culture	Growth	Line of puncture	Liquefaction		
			Type	Begins	Complete
				days	days
160.1	Best at top	Fil.-beaded	Strati.-infund.....	2	22
170.3	Best at top	Fil.-beaded	Stratiform.....	1	22
180.2	Best at top	Fil.-beaded	Napi.-strati.....	1	22
183.2	Best at top	Fil.-beaded	Napi.-strati.....	1	22
187B.1	Best at top	Fil.-beaded	Crater.-strati.....	2	22
191.	Best at top	Filiform	None.....		
194.	Best at top	Filiform	Strati.-infund.....	1	22
195.	Best at top	Filiform	Strati.-infund.....	1	22
196.	Best at top	Fil.-beaded	Napi.-strati.....	1	22
197.	Best at top	Fil.-beaded	Crater.-strati.....	2	22
198.	Best at top	Fil.-beaded	Napi.-strati.....	1	22
201.	Best at top	Fil.-beaded	Crater.-strati.....	2	22

TABLE III—(Continued)

MILK	REACTION 20			INCUBATED AT 28° C.	
Culture]	Plain milk				Litmus milk
	Coagulation begins	Reaction		Peptonization begins	Reduction of litmus†
		6 days	18 days		
	days			days	days
160.1	2	+23†	+47†	3	7-8
170.3	2	+33	+52	3	7-8
180.2	2	+36	+46	3	7-8
188.2	2	+34	+50	3	7-8
187B.1	2	+39	+49	3	7-8
191.	None	- 8	+24	None	None
194.	2			3	7-8
195.	2	+32	+53	3	7-8
196.	2	+34	+52	3	7-8
197.	2	+29	+39	3	8-9
198.	2	+10	+35	3	12
201.	2	+35	+51	3	7-8

* None of the strains changed the color of the medium.

† Figures represent increase or loss in degrees on Fuller's scale over controls titrated on same day.

‡ The indicator had very little, if any, effect on the growth.

Cohn's solution.—No growth.

Indol production.—Negative.

A survey of the published descriptions of the potato blackleg bacillus reveals the fact that half of the authors conclude that this organism produces indol, while the others arrived at an opposite conclusion. Special pains were therefore taken by the writer to determine this point accurately.

Kligler ('14), Lewis ('15), and others have criticized the use of the Salkowski-Kitasato method (conc. H_2SO_4 and $NaNO_2$) on account of its unreliability. Kligler ('14) states that this test should be discarded, since a red coloration is frequently obtained which is not due to indol; also because the reaction in cultures which really produce indol is not constant. Ehrlich's method of testing for indol production was used by the writer in his experiments. This test will detect indol in a dilution of 1:1,000,000. It has been shown that it is 10 times more delicate than the older test mentioned above. In order to make his tests even more searching, the writer employed both "Bacto" peptone and Witte peptone in making the Dunham's solutions used as culture media. All 12 strains of the blackleg bacillus were carefully tested, using controls inoculated with *Bacillus coli*, as well as blanks. A pink coloration did not appear in a single culture

except those in which *B. coli* was planted. Incidentally, it may be stated that the strain of *B. coli* used gave rise to a stronger color reaction in the Witte peptone solution than in the "Bacto" peptone solution.

Heating at 70° C. for a few minutes did not bring out the slightest trace of pink coloration in any of the cultures inoculated with strains of the blackleg organism. Cultures kept for 2 months were also tested, and negative results were obtained as before. In these cultures a yellowish or faint brown coloration was noticed after adding the test solution. Upon looking across the surface of the liquid against a dark background, the suggestion of a very pale wine coloration was obtained which in the absence of the positive test color might be mistaken for indol production. Possibly some of the authors who have reported that the blackleg bacillus produced indol were led astray by the appearance of similar color reactions in their test cultures. Repeated warnings have been given against accepting such colorations as a positive indication of indol. When compared with the pink color appearing in cultures of *B. coli* tested for indol, one is not likely to make an erroneous interpretation.

Nitrites from nitrates.—Nitrates reduced.

Strain No. 191 was the only one that did not reduce nitrates to nitrites. The starch-KI-sulphuric-acid test recommended by Smith ('05) was employed.

Ammonia production.—Moderate.

Standard nitrate broth was used as a culture medium. Duplicate tubes were inoculated, one of which was sealed with paraffin. When this was tested at the end of the tenth day, using Nessler's solution, a yellow coloration appeared in all tubes except controls. The color was slightly deeper in the sealed tubes.

Fermentation.—Acid and small volumes of gas from dextrose, lactose, sucrose, maltose, and mannite. No acid or gas from glycerin, dextrin, and potato starch. Moderate growth in closed arm of fermentation tubes, from which dissolved oxygen is driven off, in presence of dextrose, lactose, sucrose, maltose. No growth at first, feeble growth later in closed arm under similar conditions, in presence of glycerin, dextrin, and potato starch. Quantitative consumption of common hexose sugars and of sucrose, lactose, and maltose. No hydrolysis of dextrin and potato starch.

As was stated in an earlier paragraph, the several students of the blackleg bacillus differ in their published observations and conclusions as to the gas- and acid-producing capacity of different isolations of this parasite. This fact is clearly shown in the tabular summary presented herewith (table VI).

The results of the writer's fermentation studies are briefly summarized in table IV. The data presented here lead to the conclusion that the several strains of the organism studied (except No. 191) are similar and that all produce small quantities of gas and considerable acid from dextrose, lactose, and sucrose, but no acid or gas from glycerin or potato starch.

In the course of his studies the writer determined that some of the strains at hand produced gas and acid from fructose, galactose, and maltose as well (compare with results of quantitative experiments on carbohydrate consumption).

Among other things, it was determined that the gas-producing function of the blackleg bacillus varies, depending upon (1) oxygen relations, (2) the length of time cultivated in the presence of a sugar, (3) the composition of the culture medium, etc., more than upon the particular strain or strains of the organisms under observation. In one experiment (Exp. "A", table V) only the minutest volumes of gas collected in the closed arm of 3 of the tubes (lactose) out of a total of 90 inoculated (30 of lactose alone). In this set 1 per cent of the carbohydrate was added to Dunham's solution (the reaction was not adjusted) and distributed in fermentation tubes. The tubes were heated in streaming steam until the air dissolved in the solution had been expanded and driven out. The bubble which collected in the top of the closed arm of the tubes was carefully tilted off before the set was autoclaved. Small bubbles found in the closed arm of 2 of the tubes upon removal from the autoclave were tilted off. Invigorated cultures were used for inoculation, and this was accomplished in the usual manner as soon as the culture medium was cooled. The cultures were incubated at 27–28° C. A feeble to moderate growth developed in the closed arm of all tubes containing glucose, lactose, and sucrose, also in tubes under observation containing fructose, galactose, maltose, but not in closed arms of tubes containing glycerin, dextrin, and potato starch. Under the circumstances it may be concluded that the strains of the blackleg bacillus under observation grow anaerobically only when they obtain sufficient

TABLE IV
PHYSIOLOGY (FERMENTATION)

Culture No.	Medium containing Dunham's solution and:													
	Dextrose			Lactose		Sucrose		Glycerin		Potato starch		Controls		
	Gas	Reaction *		Gas	Reaction		Gas	Reaction		Gas	Reaction		Gas	
		1 dy.	5 dys.		1 dy.	5 dys.		1 dy.	5 dys.		1 dy.	5 dys.		
160.1.....	+	+4	+8	+	+2	+7	+	+3	+8	0	0	0	-1	0
170.3.....	+		+10	+		+6	+		+10	0	0	0	0	0
180.2.....	+		+7	+		+8	+		+9	0	0	0	0	0
183.2.....	+		+8	+		+7	+		+6	0	0	0	0	0
187B.1.....	+	+4	+10	+	+4	+7	+	+5	+9	0	-1	1	0	0
191.....	0		-8	0		-6	0		-7	0			0	0
194.....	+		+8	+		+5	+		+6	0		0	0	0
195.....	+	+3	+8	+	+1	+7	+	+3	+5	0	-2	-2	0	-1
196.....	+	+4	+12	+	+4	+8	+	+5	+11	0	-1	-1.5	0	0
197.....	+	+5	+10	+	+1	+7	+	+3	Lost	0	-1	-1	0	-0.5
198.....	+		+8	+		+9	+		+9	0		0	0	-0.5
201.....	+	+8	+10	+	+4	+6	+	+4	+9	0	-1	-1	0	0

* Reaction data represent gain (+) or loss (-) in degrees on Fuller's scale, over controls, after time intervals specified.

TABLE V
GAS PRODUCTION

Cult No.	Exp. A			Exp. B			Exp. C		
	Dex.	Lac.	Suc.	Dex.	Lac.	Suc.	Dex.	Lac.	Suc.
160 1.....	0	0	0	0	+	+	+	+	+
170 3.....	0	0	0	0	+	+	+	+	+
180 2.....	0	+	0	+	+	+	+	+	+
183 2.....	0	0	0	+	+	+	+	+	+
187B. 1.....	0	+	0	+	+	+	+	+	+
191.....	0	0	0	•	•	•	•	•	•
194.....	0	0	0	+	+	+	+	+	+
195.....	0	0	0	+	+	+	+	+	+
196.....	0	0	0	0	+	+	+	+	+
197.....	0	0	0	+	+	+	+	+	+
198.....	0	+	0	+	+	+	+	+	+
201.....	0	0	0	+	+	+	+	+	+

* No data.

TABLE VI
DATA ON CARBOHYDRATE REACTIONS

Investigator	Dextrose		Lactose		Saccharose		Glycerin		Diastatic action
	Gas	Acid	Gas	Acid	Gas	Acid	Gas	Acid	
van Hall.....	+	+	0	+	+	+	0	0	0
Appel.....	•	•	•	•	+	•	•	•	•
Smith.....	0	+	+	+	0	+	•	•	•
Harrison.....	0	+	+	+	0	+	0	+	+
Pethy. & Murphy.....	+	+	+	+	+	+	0	0	+
Morse.....	+	+	+	+	+	+	0	+	+
Paine.....	+	+	+	+	+	+	•	•	+
Shapov. & Edson.....	+	+	+	+	+	+	0	0	0

* No data.

oxygen for metabolic processes from oxygen-containing substances which they can break down. It may be further concluded that the gas production is inhibited by the lack of free O_2 .

In another set of tests (Exp. "B," table v) the same strains were employed as before, being cultivated in plain peptone water to which 1 per cent of the carbohydrate was added. The reaction was not adjusted. The Board of Health type of fermentation tube was employed, and sterilization was accomplished in the autoclave. After autoclaving, the tubes were set aside for a few days. Just before inoculation, the medium in each was thoroughly mixed by tilting the tube, but no free bubbles of air

were left in the closed arm. Inoculation was made in the usual manner, using invigorated strains. The cultures were kept at room temperature, 18–22° C. Gas production took place in all but 3 of the cultures (Exp. "B", table v) as was evidenced by the accumulation of from 1 to 8 or 9 per cent of a gas in the closed arm. Only very slight increases in the volume of the gas present occurred after the end of the first week. During the first 4 to 6 days there was just about as much growth in the closed arm of the tubes as in the open arm.

Still another test of the gas-producing capacity of the black-leg strains was made, using agar shake cultures. Small amounts of gas were produced by all the strains tested (Exp. "C", table v), as was evidenced by the development of a greater or less number of bubbles in the medium.

The appearance of gas bubbles in 1-day-old glucose agar slant cultures suggested the use of this type of test. A 0.75 per cent meat extract agar was used. The reaction of this medium was not adjusted. To this was added 1 per cent of the carbohydrate to be investigated. The medium was melted, cooled to 42° C., and inoculated with 3-mm. loops of invigorated broth cultures. The inoculum was thoroughly mixed and the medium aerated by a vigorous rolling between the hands. The cultures thus made were set aside in a vertical position to cool and were incubated at 27–28° C. In this type of culture it was found that there was little increase in the volume of gas produced later than the second day after inoculation, as evidenced by the number and size of the bubbles.

In following up this line of experimental work it was later determined that when agar shake cultures were sealed with paraffin a somewhat greater volume of gas was demonstrable, as evidenced by the increased number and size of the bubbles in the medium. What is even more significant: It was found that the cultivation of a particular strain (or strains) in the presence of a certain carbohydrate increased its powers to ferment that carbohydrate with the production of gas. In this way certain strains which at first produced gas very weakly, if at all, were "trained up" to ferment the particular carbohydrate in question more vigorously, as was evidenced by the production of larger and larger volumes of gas.

Diastatic action.—Absent.

The writer's conclusion is based upon the results of carefully conducted quantitative determinations of carbohydrate hydrolysis by certain of the blackleg strains at hand. In this phase of the work the hydrolysis of potato starch, as well as other carbohydrates, was investigated. No evidence was found of the slightest hydrolysis of potato starch by the blackleg bacillus after an interval of 6 days. The ordinary tests for diastatic action were made, using a nutrient agar starch jelly, and these yielded no evidence of diastatic action. These results are not in accord with those of most investigators (table VI).

Active acidity and titratable acidity of culture solutions.—While the greater part of the experimental work reported upon here was done in 1916 and 1917, the data presented in this section were obtained in May, 1922. Experiments were made in order to determine the production of acid, as shown by the determination of H-ion concentration, as well as by the relative amount of titratable acid produced by *Bacillus atrosepcticus* when cultivated in the presence of certain sugars. Subcultures of strain No. 196 were employed in these experiments.

To a nutrient broth made as before 1.0 per cent of the following sugars was added: (a) glucose ["Difco"], (b) sucrose [Merck's highest purity], (c) lactose [Merck's].

Cultures and methods.—Comparatively large-volume cultures were employed, 500 cc. of each of the sugar broths being placed in each of 3 Florence flasks of 1 liter capacity. Sterilization was accomplished in the autoclave at 15 pounds pressure for 15 minutes. In order to avoid the lag phase of growth (Chambers, '20) a rejuvenated broth culture 7 hours old was used for inoculating the sugar broth employed.

The H-ion concentration of each culture medium was determined at the time of inoculating the sugar broths and thereafter as indicated (table VII). The colorimetric method of Clark and Lubs ('17) was employed. The production of titratable acid was determined in the usual manner, using 0.507 NaOH and phenolphthalein, and the amount of titratable acid was expressed in degrees on Fuller's scale where "+10"=0.1 of 1 per cent normal HCl.

Buffer index.—Following the method of Brown ('21) the buffer index of the medium was determined in order to provide

additional data for comparison if need should arise. All 3 sugar broths employed were found to have a buffer index of 1.5 where "BI" is the sum of the reserve alkalinity and the reserve acidity, each value being expressed in per cent normal alkali or acid.

TABLE VII

H-ION CONCENTRATION AND TITRATABLE ACIDITY FOR STRAIN NO. 196

Medium	Cultures									
	Controls		1 day		2 days		5 days		11 days	
	P _H	Tit. acid.	P _H	Tit. acid.	P _H	Tit. acid.	P _H	Tit. acid.	P _H	Tit. acid.
Glucose.....	6.0	+16	5.4	+20	5.2	+22	5.0	+25	5.0	Not determined
Sucrose.....	6.0	+16	5.4	+21	5.0	+24	5.0	+25	5.0	Not determined
Lactose.....	6.0	+16	5.6	+20	5.2	+21	5.2	+25	5.1	Not determined

From table VII it may be noted at a glance that (1) some acidity is developed by *Bacillus atrosepticus* when grown in the presence of certain sugars, and (2) essentially equivalent amounts of acid are developed in the presence of all 3 sugars employed.

Attention is also directed to the fact that the major increase in H-ion concentration took place during the first 2 days, and it is believed (see Chambers, '20) that a true acidity of about P_H 5 inhibits further growth and reproduction of the organism.

SUMMARY OF COMPARATIVE STUDIES OF CAUSAL ORGANISMS

The salient characteristics of the several strains of the black-leg bacillus studied by the writer are brought together in compact form below. The scheme adopted is based upon the Descriptive Chart indorsed by the American Society of Bacteriologists in December, 1920. The digits used to designate the several "primary" and "secondary" characteristics of the strains in question may be translated in detail by reference to the foregoing text, and to the "brief characterization" column upon the descriptive chart above mentioned. The data below indicate the fact that there is very little, if any, difference between the organisms studied by the writer, except No. 191, which aside from being non-pathogenic to the potato, differs from the others in many important respects.

BRIEF CHARACTERIZATION

160.1	Montana strain	53*2-32120-2111-2222- *211-22-121
170.3	Montana strain	53 2-32120-2111-2222- 211-22-121
180.2	Montana strain	53 2-32120-2111-2222- 211-22-121
183.2	Montana strain	53 2-32120-2111-2222- 211-22-121
187B.1	Montana strain	53 2-32120-2111-2222- 211-22-121
191.	<i>B. phytophthorus</i> "as rec'd. from Appel."	53 -51230- 333-1122- 322-33-322
194.	Maine strain Morse's "SE"	53 2-32120- 111-2222- 211-22-121
195.	Maine strain Morse's "IIP"	53 2-32120-2111-2222- 211-22-121
196.	<i>B. solanisaprus</i> Harrison	53 2-32120-2111-2222- 211-22-121
197.	<i>B. atrosepticus</i> van Hall	53 2-32120-2111-2222- 211-22-121
198.	<i>B. melanogenes</i> Pethy. & Murphy	53 2-32120-2111-2222- 211-22-121
201.	<i>B. phytophthorus</i> Appel	53 2-32120-2111-2222- 211-22-121

While all differences, great or small, have been recorded, it is significant that these differences, except in the case of No. 191, are insufficient to appear in the brief characterization of each strain set down above. Such variations as may exist are, it seems to me, wholly insignificant and do not justify even varietal characterization. Within the limits of his comparative studies Morse ('17) reached similar conclusions. Among his cultures were some of the same strains used by the writer, namely, Nos. 194, 195, 196, 197, and 198.

RELATIONSHIPS AND NOMENCLATURE

Differences between the 4 "species" of the blackleg bacillus as originally described may be noted by referring to the original descriptions or to abstracts prepared by the writer (see p. 17 et seq.).

The index numbers brought together below assist in visualizing the small differences upon which certain of these "species" were established.

<i>B. atrosepticus</i> van Hall (van Hall, '02).....	5312-32120-2121
<i>B. phytophthorus</i> Appel (Appel, '03)	5?1?-32120-???1
<i>B. phytophthorus</i> Appel (Smith, '10).....	5312-32120-?212
<i>B. solanisaprus</i> Harrison (Harrison, '07)	5312-32120-1212
<i>B. melanogenes</i> Pethy. & Murphy (Pethybridge and Murphy, '10)	5312-32110-1111

* Omission of figure indicates item not determined.

Reference to the literature discloses the fact that Appel ('02, '02a) succeeded at an early date in isolating a bacterial organism which he believed to be the cause of the potato disease in question. The motive which prompted him to publish brief notes (Appel, '02, '02a) on the disease, and at the same time to propose a name (only) for the causal organism, must be left for conjecture. The fact remains that he (Appel, '03) did not publish a full description of either the organism or the disease until about a year after van Hall's ('02) dissertation was available to science.

Pethybridge and Murphy ('11) must have recognized that their species was very similar to van Hall's and Appel's, for in regard to "*B. phytophthorus*" and "*B. melanogenes*" they state: . . . "the two organisms, if not identical, are at any rate closely allied; and it is perhaps with some reluctance that we regard it as a distinct species and suggest the name *Bacillus melanogenes* for it." Relative to *B. atrosepticus* van Hall, these authors say that it has some points in common with their species, but differs by "occurring chiefly as isolated individuals, whereas ours is more frequently found in pairs. The former is also decidedly smaller in size, in spite of variations in both cases, and its action on milk appears to be different from that found in our case. *We were unfortunately unable to obtain a copy of the detailed character of B. atrosepticus* [italics not in original] before our own work was concluded."

Harrison ('07) states that "the symptoms of the Ontario disease somewhat resemble those described by Appel for the 'Schwarzbeinigkeit,' a disease which seems rather widespread in Germany." Further than this, the only comparisons made by him appear in a little display chart on page 592 of his paper. In this exhibit certain of the morphological and biological characteristics of "*B. phytophthorus*," "*B. atrosepticus*," and "*B. solanisaprus*" are compared. The differences brought out are, it seems to me, more apparent than real, as shown by the index numbers compiled (see p. 41).

As was pointed out in an earlier paragraph, Appel's name "*Bacillus phytophthorus*" appears as a *nomen nudum* in a brief paper published by him about two months previous to the appearance of van Hall's "*Bacillus atrosepticus*." Furthermore, van Hall's dissertation embodies the earliest published description of the potato blackleg disease which is sufficiently definite

and detailed to enable one to identify the disease ascribed to the bacterium named and described as the cause.

For a number of reasons Smith ('20) believes it best to retain Appel's name, especially as van Hall made very few inoculations under natural conditions, and further, because he says of his organism: "On artificial media the parasite loses its virulence very quickly." This contention loses weight in the light of a similar statement made by Smith on a preceding page (see Smith, '20, page 263), as follows: "It is common belief (*of German origin*) [italics not in original] that the organism loses virulence readily." The writer feels that Smith's further objections to the use of van Hall's name are largely refuted by the data and facts previously presented.

It was inevitable that questions involving nomenclature should have appeared in this paper and a serious attempt has been made to give all names most careful consideration. On the grounds of priority as well as for other reasons brought out above the writer believes that *Bacillus atrosepticus*¹ van Hall should stand.

REVISED DESCRIPTION OF *BACILLUS ATROSEPTICUS* VAN HALL

Index No. 5312-32120-2111.²

Microscopic features.—The potato blackleg bacillus is a small non-sporiferous, Gram-negative rod, having an average diameter of about 0.6μ and a length (1.5μ) slightly exceeding twice its diameter. The organism is actively motile by means of a few peritrichic flagella. No capsule is demonstrable by the ordinary methods now in use.

Physiological characteristics and cultural features.—The organism is non-chromogenic in agar, gelatin, nutrient broth, and other common media. On agar slants growth is moderately abundant. The surface is smooth and the luster is glistening. Agar colonies are small, round to somewhat irregular in form, and, under a magnification of 50 diameters, they appear to be granular in structure.

Gelatin is liquefied, the action being visible on the second day, if not before. Colonies on gelatin plates are white, round, and noticeably larger than those which develop on agar.

¹According to a recently suggested outline of bacterial classification (Winalow et al, '20) the name would become *Erwinia atrosepticus* nov. comb.

²See Descriptive Chart indorsed by Soc. of Am. Bacteriologists, Dec. 30, 1920.

Nutrient broth is promptly clouded, the medium becoming slightly turbid after a few days. The characteristic surface growth is a ring, though in some broths a light pellicle may develop.

Milk is promptly coagulated, the amount of titratable acid produced by the organism increasing steadily for several days. A slow peptonization of the curd begins on the second or third day.

In Cohn's solution there is no growth. In Uschinsky's solution the growth is copious, but the fluid does not become viscid.

No indol is demonstrable by the Ehrlich test, in either young or old cultures grown in peptone water made with (1) Witte peptone, or (2) Bacto peptone. Nitrates in nitrate broth are reduced to nitrites without the formation of gas. Ammonia production is feeble to moderate.

Carbohydrate reactions.—Acid and small volumes of gas are produced from dextrose, galactose, sucrose, maltose, and mannite. The gas-producing capacity is not particularly characteristic. No acid and no gas are produced from glycerin, dextrin, and potato starch. Quantitative consumption of the common hexose sugars, glucose, fructose, and galactose, has been demonstrated. Likewise, the organism consumes sucrose, lactose, and maltose. A sugar concentration of 0.25 per cent is ample. Diastatic action is absent, both in respect to starch and dextrin.

Enzymes.—This organism secretes several carbohydrate enzymes, as is shown by its action on as many different saccharides; cytase also is probably produced.

The optimum temperature for growth is about 26° C., with no growth at 37.5° C. The organism withstands considerable extremes of cold, being found viable in soil cultures exposed for 24 hours to temperatures ranging from -6.7° to -28.2° C.

The blackleg bacillus is quite resistant to drying and remains viable for long periods of time on plain beef agar.

Virulent strains produce a necrosis of the stem and tubers of the potato. Parenchymatous tissue is almost exclusively affected, and a blackening of the diseased tissues is characteristic.

III. CARBOHYDRATE UTILIZATION BY STRAINS OF THE BLACKLEG BACILLUS AND OTHER MICROORGANISMS

In spite of the fact that the strains of the blackleg bacillus employed by the writer in earlier studies showed no signs of diastatic action on potato starch jelly, it was nevertheless somewhat difficult to conceive of a virulent parasite of the potato totally lacking the power of hydrolyzing starch. This thought, together with a rather extensive knowledge of the gas- and acid-producing capacity of this bacillus, suggested the desirability of investigating these relations in a quantitative manner. The writer had employed Shaffer's ('14) modification of the volumetric method of Bertrand for the quantitative estimation of reducing sugars in the presence of proteins and albumins. Familiarity with this method led me to conclude that with its aid reliable data could be obtained concerning the quantitative consumption of certain sugars and starch by the blackleg bacillus as well as by other microorganisms with which it appeared. Very little, if anything, had been published upon the quantitative consumption of carbohydrates. Accordingly, plans were made to investigate quantitatively these relations in certain of the strains of the potato blackleg bacillus as well as in other species of bacteria. Some points of interest in this connection have, however, recently been contributed by Besson, Ranque, and Senez ('19). Working with *Bacillus coli* in nutrient broth containing varying amounts of dextrose, they found that when less than 0.4 per cent dextrose was furnished all the sugar was removed in 24 hours.

MATERIALS AND METHODS

After numerous experiments a method was developed whereby reliable data were obtained. The work of Shaffer and Hartmann ('21) on the iodometric determination of copper and its use in sugar analysis had not appeared at the time the writer performed the experiments reported herewith. Undoubtedly, the newer methods would have greatly facilitated prosecution of this phase of the work, but it is doubtful if more accurate results could have been obtained.

Certain strains of the blackleg bacillus used in the preceding studies were selected for investigation and comparison in this connection, i. e., Nos. 160.1, 180.2, 183.2, 187B.1, 195, 197, 198, 201. Pure cultures of the following were also tested: *Bacillus coli*

communis, *B. vulgaris*, and a species of yeast isolated from a compressed yeast cake.

The basic culture solution used throughout was a Dunham's solution to which 1 per cent of the following test substances (Merck's, except starch) was added: (1) glucose (H. P.), (2) fructose (cr.), (3) galactose (powder), (4) saccharose (H. P.), (5) lactose (H. P. cr.), (6) maltose (powder), (7) dextrin (H. P. powder), and (8) potato starch (Heil's).

The nutrient solutions used were not adjusted by the addition of acid or alkali. As a matter of fact, it was determined in a preliminary experiment that if small amounts of acid (HCl) were added to adjust the reaction a considerable hydrolysis of certain disaccharide sugars resulted during the process of sterilization, regardless of whether the latter was accomplished in the autoclave or in the Arnold sterilizer (table ix). The Dunham's solution employed as a basis of the nutrient solutions used was found to be P_H 7.0, approximately. The P_H of the several sugar broths employed was not determined, but it is unlikely that they differed markedly in H-ion concentration from the Dunham's solution.

Invigorated cultures of the several microorganisms at hand were employed. Ten-cc. volumes of the media were measured into special culture tubes, called wasp tubes (see pl. 2, fig. 5). (The wasp tube was suggested to me by Dr. G. W. Freiberg, formerly of this laboratory, and has greatly facilitated and expedited my work.) These were plugged in the usual manner and sterilization was accomplished in the autoclave (see table ix). Duplicate cultures were made, and a number of controls were carried along with each series of cultures. All cultures were incubated at 28° C. for 6 days, as it was found that there was no point in carrying the cultures for a longer period. Chambers ('20) has recently contributed a convincing explanation of why evidences of the metabolic activities fail after a few days in certain sugar-broth cultures.

Shaffer's ('14) method for the quantitative estimation of reducing sugars was used with slight modifications. This development of the permanganate titration method of Bertrand is of special value to workers in plant physiology because it enables one quickly and accurately to determine the amount of sugar as low as 2 mgms. The copper-reducing value of the culture

medium in which the microorganisms in question were grown was determined by this method, as was the copper-reducing value of control portions. Differences were carefully noted and carbohydrate utilization estimated by reference to Shaffer's table of copper-glucose equivalents, having first determined by calculation that 1 cc. of the N/20 permanganate is equivalent to 3.18 mgms. copper. For the technique of the complete method see under "Technique."

According to Plimmer ('15) pure lactose reduces Fehling's solution 71 per cent as strongly as glucose; pure maltose 64 per cent as strongly as glucose. These figures were employed in making estimates of the amounts of these disaccharides consumed by the microorganisms under consideration. It was assumed in this work that the fructose and galactose used reduced Fehling's quite as strongly as glucose. Certain dextrans reduce Fehling's slightly, but the product which I employed did not; nor did the saccharose and potato starch. Where it became necessary to hydrolyze these last-named substances in order to determine whether there had been utilization of the substance as such, the following methods were used:

Sucrose.—Davis and Daish ('13) employed citric acid in preference to a mineral acid for inverting cane sugar. In dealing with plant extracts, in which there was an accumulation of sodium acetate, they employed 10 per cent citric acid.

Method.—(1) Make solution faintly acid to methyl orange by addition of a few drops of concentrated H_2SO_4 ; (2) add 5 per cent by weight of citric acid; (3) autoclave for 15 minutes at 15 pounds pressure, cool, and neutralize to phenolphthalein with NaOH.

Starch and dextrin. *Method.*—(1) To a 1 per cent starch suspension (10 cc.) add 0.3–0.5 cc. conc. HCl and 10 cc. distilled water; (2) heat in autoclave for 15 minutes at 15 pounds pressure; (3) cool, neutralize to phenolphthalein with NaOH, and estimate as glucose. The amount of glucose times 0.9 equals the weight of starch hydrolyzed.

Maltose and lactose. *Method.*—(1) To 10 cc. of a 1 per cent solution of the sugar add 0.3–0.5 cc. conc. HCl (sp. gr. 1.16) and 10–15 cc. distilled water; (2) heat in autoclave for 15 minutes at 15 pounds, cool and neutralize to phenolphthalein by addition of NaOH; (3) dilute and estimate as glucose.

Complete reduction of these sugars is difficult, if not impossible, to accomplish by boiling for 60 to 90 minutes. Ordinarily, inversion is complete to the extent of 96-97 per cent. Heating for a longer time will accomplish more inversion, but is offset by an increasing destruction of the resulting monosaccharides.

A trial of the method developed by Ling and Rendle ('05) showed that this could not be used because of (1) a clouding of the culture solution immediately upon addition of the Fehling's, making it practically impossible to observe discharge of blue color upon nearing the end point, and (2) the CuO_2 does not settle readily. The trouble is probably due largely to the amino acids of the peptone in the Dunham's solution employed (cf. Davis and Daish, '13).

REAGENTS AND SOLUTIONS

Fehling's solution.—It was found best to use the Soxhlet-Fehling solution, preliminary tests having demonstrated that the solution as modified by Allihn contained too much alkali, when made up with NaOH at the rate of 178 gms. per liter.

Cuprous oxide solvent.—The ferric sulphate-sulphuric acid-cuprous oxide solvent recommended by Shaffer ('14) was employed. This solvent contains 10 per cent ferric sulphate, $\text{Fe}_2(\text{SO}_4)_3$, in 25 per cent sulphuric acid. It is a very active solvent, and care must be taken in watching for the end point. To be efficacious it should be made up by mixing equal parts of a 20 per cent aqueous ferric sulphate and 50 per cent sulphuric acid. The ferric sulphate should be dissolved in hot water, filtered while hot, and the warm acid mixed with the warm ferric sulphate solution. Just enough permanganate should be added to oxidize any ferrous salt which may be present.

Potassium permanganate solution.—The potassium permanganate used was made up according to Olsen ('10). Freshly prepared and carefully standardized N/20 permanganate was employed for titration. One cc. of N/20 permanganate is equivalent to 3.18 mgms. of copper.

TECHNIQUE

The determination of reducing sugars.—The following technique was followed in determining amounts of reducing sugars in carbohydrate broth cultures of bacteria. Certain slight modi-

fications of Shaffer's ('13) original method appear, but these are largely omissions of certain steps which were determined by preliminary experimentation to be unnecessary. For instance, acetic acid and colloidal iron were not used in the process because it was determined that the small amounts of albuminous material present in the cultures did not interfere with the determination of the reducing sugars present. The precautions emphasized by Shaffer were observed throughout. The cuprous oxide was titrated immediately upon being dissolved and without removal from the centrifuge tube in which it was precipitated and thrown down. In order to avoid breakage of the centrifuge tubes when these were plunged in the water bath circular wire baskets (pl. 2, fig. 5) having wooden bottoms and tops, with holes for 2 centrifuge tubes, were used. No oxidizable substance other than the sugar was allowed to get into the solution, and the cuprous oxide solvent employed was made from chemically pure substances in order to avoid presence of ferrous iron. This was further assured by adding a trace of permanganate.

Procedure.—The procedure outlined in the steps given below was followed in carrying out all experiments in this phase of the work:

(1) A large volume of Dunham's solution was made up, autoclaved, and then filtered through paper.

(2) To separate portions of this solution, 1 per cent of the test substances (see list p. 46) was added and dissolved.

(3) Exactly 10 cc. of each of the 8 nutrient solutions thus prepared were placed in each of 10 special culture tubes ("wasp" tubes, pl. 2, fig. 5). These tubes were large enough to permit dilution of the culture to exactly 60 cc. This was made possible by carefully standardizing each tube and permanently etching the 60-cc. mark on the slender neck. Had these tubes been drawn out and graduated to hold 50 cc., calculations would have been simplified.

(4) The tubes were plugged with cotton in the usual manner and the contents sterilized in the autoclave by exposing for 15 minutes at 15 pounds pressure. One of the preliminary experiments performed showed that this method, if carefully controlled, did not hydrolyze the di- and polysaccharides used (table ix).

(5) Each of the 8 media was inoculated with a loopful of an invigorated culture of the organism to be tested for its ability to hydrolyze the test substances employed. Two uninoculated "control" tubes of each sugar broth were carried along, incubated, and tested in identically the same manner as the inoculated cultures in each set.

(6) The sets thus made ready were incubated at 27–28°C. for 6 days, preliminary tests having shown that there was no point in incubating the cultures for more than 5 or 6 days (cf. Chambers, '20).

(7) The culture medium in each "wasp" tube was next diluted 6 times by adding distilled water to the 60-cc. mark. The contents of each was thoroughly mixed.

(8) Next, exactly 10 cc. of the diluted culture medium were removed and placed in a 50-cc. lipped centrifuge tube (pl. 2, fig. 5), which was marked to correspond to the culture tube from which the medium came. These 10-cc. volumes, then, represented 1/6 of the original culture solution, i.e. 1.666 cc. of the original 10; and if the 10 cc. of the culture solution contained 0.1 gm. of a carbohydrate a 10-cc. sample from the diluted culture should contain 0.0166 gm. of the substance provided none had been consumed.

Each tube in the set was handled in exactly the same way, the same pipette being used throughout.

(9) Ten cc. of freshly mixed Soxhlet-Fehling's solution were promptly added to the 10-cc. volumes in the centrifuge tubes.

(10) All were quickly placed in the special baskets and immersed in a water bath (pl. 2, fig. 5) where they were exposed at the boiling point for exactly 10 minutes.

(11) Upon removal from the bath the tubes were nearly filled with distilled water, pairs balanced, and centrifuged¹ at a moderate speed for 3–4 minutes.

(12) Upon removal from the centrifuge the Fehling's was cautiously decanted over a white dish, the cuprous oxide at the bottom being disturbed as little as possible, for fear of loss.

(13) This having been done each tube was again filled with distilled water in order to wash the precipitate, pairs balanced, and again centrifuged for about 4 minutes.

(14) When finally removed from the centrifuge the wash water was decanted as completely as possible without disturbing the cuprous oxide in the bottom (if any was present).

(15) The cuprous oxide was then promptly dissolved in as small a volume of the ferrous sulphate-sulphuric acid solvent as possible.

(16) The copper was then titrated immediately against N/20 potassium permanganate. No attempt was made to remove the dissolved cuprous oxide from the centrifuge tube, the actual amount of copper present being determined by titrating directly into the tube.

(17) Using the figures thus obtained and remembering that the 10-cc. sample titrated represents 1.666 cc. of the original culture medium,

¹ The No. 1 centrifuge of the International Instrument Co. was used.

the actual amount of sugar (as glucose) left was calculated by reference to Shaffer's ('14) table of copper-glucose values. In order to determine consumption of the non-reducing carbohydrates in use, i.e., sucrose, dextrin, and starch, it was necessary to accomplish inversion by heating in the presence of an acid (see "Methods," p. 47, et. seq). This having been accomplished, the procedure was exactly as outlined above.

The control tubes containing the several carbohydrate broths were handled in exactly the same manner as the cultures, except for inoculation, and were a component of each experiment.

EXPERIMENTAL DATA

The data relative to carbohydrate utilization are presented in tables IX-XIII. Where practicable the amounts of the test substance consumed are given in mgms. of the carbohydrate supplied. The most complete data are, for obvious reasons, given in terms of mgms. of glucose. The writer believes that the figures are reliable to the extent of showing relative differences in carbohydrate utilization as well as the amounts of the test substances consumed by each of the several species and strains of the microorganisms employed.

A slight error in the sugar data presented may have developed, due to the fact that it was necessary to make interpolations between figures available in tables IX-XIII, in order to determine the sugar equivalent of the copper values obtained from my own permanganate titration data. The following figures and statements are presented to illustrate the methods and the calculations employed in obtaining the sugar values presented in the above-mentioned tables. The specific case of *Bacillus coli* in the presence of glucose is selected for this illustration. The 10-cc. sample of the culture taken for treatment with Fehling's and subsequent titration represented (as in all other cases) 1.666 cc. of the original 10-cc. sugar broth in which the organism was cultivated. It was not necessary to account for possible loss due to evaporation, since the cultures were made up to 60 cc. before sampling.

Having determined the number of cc. of permanganate for 1 cc. of the original culture solution as 4.15, then the number of mgms. of copper is represented by 13.2, determined by multiplying 4.15 by the factor 3.18 (it was previously determined that 1 cc. of N/20 permanganate is equivalent to 3.18 mgms. copper).

Next, the glucose equivalent of the 13.2 mgms. copper is estimated by interpolating in Shaffer's table of copper-glucose equivalents, and finally, the amount of glucose consumed or used by the organism is found by determining the difference between the amounts of glucose recovered in the control and in the culture.

Where amounts of sucrose are given they were determined as follows:

$$96 : 100 :: A : X,$$

where the reducing ratio

$$\frac{\text{Glucose}}{\text{Invert sugar}} = 0.96,$$

and A is the invert sugar ("glucose") value obtained by titration and calculation, and X equals the true glucose value.

Then:

$$360 : 342 :: X : Y,$$

where 360 is the molecular weight of 2 molecules of $C_6H_{12}O_6$ and 342 is the molecular weight of $C_{12}H_{22}O_{11}$. Then, where X equals true glucose value determined above, Y equals sucrose consumed. Fructose and galactose consumption were calculated from the "glucose" values obtained, using the reducing ratios of these sugars as given by Browne ('12):

$$\text{Ratio } \frac{\text{Glucose}}{\text{Fructose}} = 0.92,$$

$$\text{Ratio } \frac{\text{Glucose}}{\text{Galactose}} = 0.90.$$

Consumption of lactose and maltose is directly determinable, since both these sugars are reducing sugars. Consumption of lactose, maltose, dextrin, and starch, as glucose, was determined by hydrolyzing portions of both control and culture solutions (see "Methods," p. 47). For well-known reasons no attempt was made to calculate consumption of dextrin and starch as such, nor was it thought to be worth while to convert the galactose-glucose value obtained by titrating the hydrolyzed lactose into glucose. It should be borne in mind that the glucose values given for dextrin and starch were derived by multiplying the glucose value obtained by titrating the hydrolyzed product by 0.9. Without doubt the figures in the tables representing consump-

tion are slightly erroneous, since, as is well known, more or less destruction of hexose sugar takes place in the process of hydrolyzing carbohydrates by heating in the presence of an acid. It may be recalled, too, that variability in the reducing power of lactose and of maltose is a characteristic of these disaccharides. According to Browne ('12, p. 402), succeeding portions of lactose and maltose will vary in reducing power according to the amount of free alkali, time of boiling, etc. "This peculiarity of maltose and lactose" he says, "is explained by a slight hydrolysis of the sugar into monosaccharides of higher reducing power" during the process of heating. Browne believes that a slight inversion of this kind takes place to a greater or less extent with all higher saccharides (including sucrose) upon boiling with Fehling's solution.

These facts may account in part for the inconsistency of findings reported in the literature, to show that moist heat of moderate degree causes hydrolysis of certain higher disaccharides and other carbohydrates, such as dextrin and starch.

In carrying out a number of tests preliminary to the major work reported upon in connection with this phase of the investigation, one experiment is worthy of particular consideration,—that made to determine the effect of moist heat upon certain carbohydrates. The temperatures used were those encountered in sterilizing solutions in the autoclave and by the discontinuous method. Three different carbohydrate-containing solutions were made up: (1) distilled water with 0.5 per cent of the carbohydrates to be tested, (2) Dunham's solutions, acidulated by the addition of 0.4 per cent N/1 HCl, separate portions of which contained 1 per cent of the carbohydrates, (3) a plain Dunham's solution which contained 1 per cent of the carbohydrates being tested. Each of these lots was separated in 3 portions. The controls were not heated and were titrated immediately. The second set was autoclaved at 15 pounds (121.3° C.) for 15 minutes, the total length of time above 100° C. being 45 minutes. The third set was exposed in an Arnold steamer for intervals of 20 minutes at about 99.5° C. on 3 consecutive days. The findings appear in table ix, and it will be seen from the tabulated data that sterilization of certain saccharide broths was accomplished by either method without significant hydrolysis of the carbohydrates present when no acid (HCl) was added. Certain of the monosaccharides, as well as the disaccharides lac-

tose and maltose, dissolved in the Dunham's solution appeared to have been slightly altered during the process of sterilization in the Arnold sterilizer, but it seems doubtful if the changes which are indicated by the figures are significant even in the case of lactose and maltose, since, as has been pointed out, these sugars are characterized to a certain extent by a variability in reducing power.

On the other hand, reference to table ix emphasizes clearly the fact that a considerable hydrolysis of the di- and polysaccharides occurred in the acidulated broths exposed in the autoclave, and to a lesser extent in those exposed in the steamer. It also appears that sucrose is most markedly affected. According to my tests, lactose, maltose, and potato starch in slightly acidulated broth are not hydrolyzed by exposure to moist heat in the process of sterilization at 100° C.

The data given below are of interest when compared with the quantitative data presented in table ix. These data, furthermore, serve as a check on the other type of analysis.

TABLE VIII
RESULTS OF COLORIMETRIC TEST WITH IODINE.

Carbohydrate	Controls	Autoclave	Arnold Steamer
Dextrin	Magenta	No color	Very pale magenta
Potato starch	Blue	Magenta	Blue

In the other tables (ix-xiii) are presented data bearing on the quantitative consumption of chosen carbohydrates by a few microorganisms. Nine of the 12 cultures employed were representative strains of the blackleg bacillus and among the 9 were the 4 "species" of this bacillus, which, together with the other strains, were studied comparatively at an earlier date (see Part II). In addition to these, cultures of *Bacillus coli*, *B. vulgaris*, and a species of yeast (probably a species of *Saccharomyces*) were investigated. For the sake of facilitating comparisons the amounts of carbohydrate consumed, in terms of glucose, are assembled in table xiv.

DISCUSSION

The data presented in tables ix-xiv call for very little further commentary, but the action of strain numbers 195 and 196 (table xi and xii) in the presence of sucrose attract special

attention as being extraordinary for the bacillus represented. It would appear that these strains, unlike others of *Bacillus atrosepticus*, inverted cane sugar faster than the hexose by-products were used up. Also, it will be noted that these strains consumed nearly triple the average amount of sucrose used by the other blackleg strains. In this respect these cultures are comparable with *B. vulgatus*, which is the only other one of the 12 which consumes sucrose so strongly and, in addition, inverts this saccharide faster than it uses up the invert sugar. It is possible that these 2 cultures (Nos. 195 and 196) were contaminated during the course of these experiments. Neither attracted attention because of extraordinary cultural features at the time and so were not plated out.

Figures are presented in tables XI, XIII, and XIV which indicate a very slight consumption of dextrin by the blackleg strains number 187B.1 and 198. Also, it would appear that minute quantities of starch were used by strain No. 201 (table x), and it would seem that lactose was attacked by *B. vulgatus* as well as maltose by the yeast species (tables XIII and XIV). The writer is of the opinion, however, that these data in themselves are slightly misleading. It seems to him more than likely that they appear as the result of small discrepancies which are likely to develop in work of this nature and for reasons which have been stated above. It will be noted that the strain of *Bacillus coli* investigated did not hydrolyze sucrose nor consume it. *Bacillus vulgatus* as cultivated in these experiments did not utilize lactose nor did it use more than about a third as much galactose as glucose and fructose. Both these bacteria, it will be noted, attack with considerable avidity all the other carbohydrates employed. The yeast species used was unable to attack and use carbohydrates presented other than glucose and fructose. These sugars, however, were totally eliminated from the broth by this microorganism.

The several strains of *Bacillus atrosepticus* employed in these tests were quite alike in respect to their attack upon and use of the several carbohydrates presented. It appears likely that 15-20 per cent of the amount of the carbohydrates presented would have been sufficient to supply the strains employed, under the conditions of the experiments. The strains of the blackleg bacillus, furthermore, utilized sucrose, leaving no invert sugar behind except in the cases cited above of strains Nos. 195 and

TABLE IX

EFFECT OF METHOD OF STERILIZATION ON CARBOHYDRATES

Carbohydrates tested	0.5% in aqueous sol.						1% in Dunham's sol. +10 by HCl						1% in Dunham's sol. Not adjusted					
	Contrl.*		Autoclave		Arnold		Contrl.		Autoclave		Arnold		Contrl.		Autoclave		Arnold	
	Total	Gain	Total	Gain	Total	Gain	Total	Gain	Total	Gain	Total	Gain	Total	Gain	Total	Gain	Total	Gain
	recov- ery	due to heat	recov- ery	due to heat	recov- ery	due to heat	recov- ery	due to heat	recov- ery	due to heat	recov- ery	due to heat	recov- ery	due to heat	recov- ery	due to heat	recov- ery	due to heat
	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.
Glucose.....	4.7	4.7	0	4.7	0	4.7	10.2	10.2	0	10.2	0	10.2	0	10.6	10.5	0	10.6	0
Fructose.....	4.1	4.1	0	4.1	0	4.1	9.07	9.14	0.07	9.14	0.07	9.14	0.07	9.1	8.95	0	9.17	0.07
Galactose.....	4.4	4.1	0	4.1	0	4.1	9.26	9.25	0	9.27	0.01	9.27	0.01	9.15	9.2	0	9.2	0.05
Sucrose.....	0	0	0	0	0	0	0	11.5 (11.0)	11.5	11.3 (10.8)	11.3	11.3	0	0	0	0	0	0
Lactose.....	3.3	2.8	0	3.3	0	3.3	6.35	7.27	0.92	6.35	0	6.35	0	5.35	5.44	0.09	5.5	0.2
Maltose.....	2.32	2.31	0	2.31	0	2.31	4.6	6.07	2.67	4.55	0	4.55	0	3.9	3.89	0	4.18	0.28
Dextrin.....	0	0	0	0	0	0	0	1.44	1.3	0.85	0.76	0.76	0	0	0	0	0	0
Potato starch.....	0	0	0	0	0	0	0	1.04	0.9	0	0	0	0	0	0	0	0	0

* Not heated.

TABLE X
CARBOHYDRATE CONSUMPTION OF *BACILLUS ATROSEPTICUS* STRAINS

	Controls				No. 201				No. 197			
	Total recovery		Total recovery		Total recovery		Consumption		Total recovery		Consumption	
	Carb.*	Gluc.†	Carb.	Gluc.	Carb.	Gluc.	Carb.	Gluc.	Carb.	Gluc.	Carb.	Gluc.
	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.
Glucose.....	9.3	9.3	8.2	8.2	1.1	1.1	8.4	8.4	0.9	0.9	0.9	0.9
Fructose.....	8.4	9.2	7.3	8.0	1.1	1.2	7.4	8.0	1.0	1.2	1.2	1.2
Galactose.....	8.8	9.8	7.7	8.6	1.1	1.2	7.6	8.5	1.2	1.3	1.3	1.3
Sucrose.....	0	9.8	0	8.5	1.2	1.3	0	8.7	1.0	1.1	1.1	1.1
Lactose.....	5.8	(9.4)†	4.7	(8.2)†	1.1	1.9	4.7	(8.4)†	1.1	2.0	1.1	2.0
Maltose.....	4.8	9.4	4.0	8.6	0.8	0.8	4.1	7.0	0.7	0.8	0.7	0.8
Dextrin.....	0	8.1	0	8.1	0	0	0	8.1	0	0	0	0
Potato starch.....	0	8.2	0	8.1	0	0.1	0	9.0	0	0	0	0

* Carbohydrate.

† Glucose.

‡ Invert sugar value in parenthesis.

TABLE XI
CARBOHYDRATE CONSUMPTION OF *BACILLUS COLI* AND *B. ATROSEPTICUS* STRAINS

	Controls				<i>Bacillus coli</i>						<i>B. atrosepticus</i>					
	Total recovery		Gluc.		Total recovery		Consump- tion				Total recovery		Consump- tion			
	Carb.	mgn.	Carb.	mgn.	Carb.	mgn.	Carb.	mgn.	Carb.	mgn.	Carb.	mgn.	Carb.	mgn.	Carb.	mgn.
Glucose.....	9.5	9.5			6.0	6.0	3.5	3.5			8.3	8.3	1.2	1.2	7.8	7.8
Fructose.....	8.5	9.2			5.5	6.0	3.0	3.2			7.7	8.3	0.8	0.9	7.5	8.1
Galactose.....	8.7	9.7			5.4	6.0	3.3	3.7			7.8	8.7	0.9	1.0	7.3	8.1
Sucrose.....	0	9.7 (9.3)			0	9.7 (9.3)	0	0			0	8.7 (8.3)	0.9	1.0 (2.1)	2.2* (5.8)	6.0* (5.8)
Lactose.....	5.7	9.0			3.3	6.3	2.4	2.7			5.1	7.8	0.6	1.2	4.9	7.8
Maltose.....	4.5	9.4			3.2	6.2	1.3	3.2			4.2	8.7	0.3	0.7	3.2	7.0
Dextrin.....	0	8.2			0	6.8	+	1.4			0	8.0	0	0.2	0	8.2
Potato starch.....	0	8.1			0	6.0	+	2.1			0	8.1	0	0	0	8.2

* These figures seem to indicate an extraordinary sucrose consumption (but see pages 55 and 62). Certain of them indicate that the sucrose was inverted faster than used by the organism. See also No. 196, table xii.

TABLE XII

CARBOHYDRATE CONSUMPTION BY *BACILLUS ATROSEPTICUS*

	Controls				No. 195				No. 180.2				No. 180.1			
	Total recovery		Consumption		Total recovery		Consumption		Total recovery		Consumption		Total recovery		Consumption	
	Carb.	Gluc.	Carb.	Gluc.	Carb.	Gluc.	Carb.	Gluc.	Carb.	Gluc.	Carb.	Gluc.	Carb.	Gluc.	Carb.	Gluc.
	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.	mgms.
Glucose.....	9.5	9.5	8.7	8.7	0.8	0.8	8.2	8.2	1.3	1.3	8.4	8.4	8.4	8.4	1.1	1.1
Fructose.....	8.7	9.5	7.6	8.3	1.1	1.2	7.4	8.0	1.3	1.5	7.7	8.4	7.7	8.4	1.0	1.1
Galactose.....	8.7	9.7	7.4	8.2	1.3	1.5	7.4	8.2	1.3	1.5	8.2	9.1	8.2	9.1	0.5	0.6
Sucrose.....	0	9.7 (9.3)	2.0*	6.1 (5.9)	3.4	3.6*	0	7.5 (7.2)	2.1	2.2	0	8.7 (8.4)	0	8.7 (8.4)	0.9	1.0
Lactose.....	6.0	9.5	5.0	8.0	1.0	1.5	4.9	7.4	1.1	2.1	5.6	7.8	5.6	7.8	0.4	1.7
Maltose.....	4.5	9.2	4.4	8.8	0.1	0.4	4.0	8.0	0.5	1.2	3.9	8.6	3.9	8.6	0.6	0.6
Dextrin.....	0	8.1	0	8.1	0	0	0	8.1	0	0	0	8.1	0	8.1	0	0
Potato starch.....	0	8.1	0	8.1	0	0	0	8.1	0	0	0	8.1	0	8.1	0	0

* Cf. also No. 196 table XI.

TABLE XIII

CARBOHYDRATE CONSUMPTION BY *BACILLUS VULGATUS*, A YEAST, AND STRAINS OF *B. ATROSEPTICUS*

	Controls			<i>Bacillus atrosepticus</i>						<i>Bacillus vulgatus</i>						A yeast					
	Total recovery			No. 198			Total recovery			No. 183.2			Total recovery			Total recovery			Total recovery		
	Carb.	Gluc.	mgms.	Carb.	Gluc.	mgms.	Carb.	Gluc.	mgms.	Carb.	Gluc.	mgms.	Carb.	Gluc.	mgms.	Carb.	Gluc.	mgms.	Carb.	Gluc.	mgms.
Glucose.....	9.4	9.4	8.4	8.4	1.0	1.0	8.2	8.2	1.2	1.2	1.2	1.2	5.4	5.4	4.0	0	0	0	All	All	All
Fructose.....	8.6	9.4	7.4	8.0	1.2	1.3	7.2	7.8	1.4	1.6	1.6	1.6	4.6	5.0	4.0	0	0	0	All	All	All
Galactose.....	8.7	9.7	7.7	8.5	1.0	1.2	7.8	8.6	0.9	1.1	7.4	8.2	1.3	1.5	1.5	9.1	10.0	0	0	0	0
Sucrose.....	0	9.6	0	8.3	1.2	1.3	0	8.2	1.3	1.4	6.0†	7.3	2.2	2.3	0	9.6	0	0	0	0	0
Lactose.....	5.9	9.0	5.0	7.4	.9	1.6	4.9	6.9	1.0	2.1	5.8	8.9	0.1	0.1	0.1	6.0	9.2	0	0	0	0
Maltose.....	4.5	9.3	4.9*	7.4†	?	1.9	4.1	8.5	0.4	.8	2.3	5.9	2.2	3.4	4.5	9.1	0	0.2	0	0	0
Dextrin.....	0	8.2	0	8.0	0	0.2	0	8.2	0	0	1.3†	4.9	0	3.3	0	8.2	0	0	0	0	0
Potato starch.....	0	8.1	0	8.2	0	0	0	8.4	0	0	1.0†	4.5	0	3.7	0	8.2	0	0	0	0	0

* Clumps of growth in culture.

† This organism evidently hydrolyzes certain carbohydrates, splitting off invert sugar faster than it consumes it.

TABLE XIV

AMOUNTS OF CARBOHYDRATE CONSUMED BY CERTAIN MICRO-ORGANISMS

Test substance	Mgms. of glucose											
	B. coli	B. vulgatus	A yeast	Bacillus atrosepticus								
				201	198	197	196	195	187B.1	183.2	180.2	160.1
Glucose.....	3.5	4.0	9.4	1.1	1.1	0.9	1.7	0.8	1.2	1.2	1.3	1.1
Fructose.....	3.2	4.4	9.4	1.2	1.3	1.2	1.1	1.2	0.9	1.6	1.5	1.1
Galactose.....	3.7	1.5	0	1.2	1.2	1.3	1.1	1.5	1.0	1.1	1.5	0.6
Sucrose.....	0	2.3	0	1.3	1.3	1.1	3.7*	3.6*	1.0	1.4	2.2	1.0
Lactose.....	2.7	0.1	0	1.9	1.6	2.0	1.2	1.5	1.2	2.1	2.1	1.7
Maltose.....	3.2	3.4	0.2	0.8	1.8	0.8	1.4	0.4	0.7	0.8	1.2	0.6
Dextrin.....	1.4	3.3	0	0	0.2	0	0	0	0.2	0	0	0
Potato starch.....	2.1	3.7	0	0.1	0	0	0	0	0	0	0	0

* Probably too high. See pages 55, 62.

196. *Bacillus coli* and *B. vulgatus* were the only organisms investigated that attacked and used dextrin and starch. The latter, however, appears to have been able to break down both of these substances more rapidly than it utilized the derived sugars (table XIII).

In respect to the amounts of carbohydrates actually utilized, it will be noted upon reference to table XIV that with the several microorganisms under observation (excepting the yeast) the amount is relatively small as compared with the total amount supplied (10 mgms. per cc.). The single instance of *Bacillus vulgatus* on fructose is the only one where the actual amount consumed approaches 50 per cent of the total amount of the sugar provided. The different strains of *Bacillus atrosepticus* consumed only about 10 to 20 per cent of the available supply of carbohydrates. *Bacillus coli* utilized about 30 per cent, and *B. vulgatus* about 40 per cent of the supply. In the light of these findings it would appear that the usual recommendation relative to the amount of carbohydrate that should be supplied in a nutrient medium was a most liberal one. *Bacillus coli* and *B. vulgatus* are usually thought of as being particularly active in fermenting certain saccharides, especially when compared to most plant pathogens. As it is possible that there are numerous bacterial species for which the optimum sugar concentration is 1 per cent or even more, it is doubtful if a less liberal recommendation should be made until the whole matter has been more extensively investigated. However, in the light of these findings, as well as those of Besson, Ranque and Senez ('19), Chambers ('20), Wolf and Foster ('21), and others, it appears that *B. coli*, also some plant pathogenic bacteria, require a less amount of sugar than 1 per cent for optimum development. Besides, there is abundant evidence which indicates that most plant pathogenic bacteria require a very much lower sugar concentration than 1 per cent (cf. also sugar consumption by *B. atrosepticus*, table XIV).

Further evidence of the similarity of the several strains of the blackleg bacillus employed throughout these investigations is brought out in tables X-XIII.

GENERAL SUMMARY

I

The blackleg disease of Irish potatoes in North America and Europe is caused by a Schizomycete which should bear the name *Bacillus atrosepticus* van Hall.

The following names are to be considered only as synonyms: *Bacillus phytophthorus* Appel, *B. solanisaprus* Harrison, *B. melanogenes* Pethybridge & Murphy.

The index number "5312-32120-2111" very briefly describes *Bacillus atrosepticus* van Hall, the number being based on the results of the writer's comparative studies.¹ A revised description of this organism appears on pages 43-44.

The blackleg parasite grows best at about 26° C. It withstands extremely low temperatures (-28.2°C.) for several hours.

This organism produces acid and a small volume of gas from each of a number of sugars. The gas-producing capacity is relatively weak, but this capacity can be built up to a certain extent by constant cultivation in the presence of the sugars which it is able to utilize.

The pathogen infects the vines and the tubers of the potato.

Virulence of the parasite, as tested by artificial inoculation, appears to be dependent upon a rather delicate balance of temperature and water relations, and upon the sugar content of the tissues inoculated.

The "incubation period" varied in the experiments under observation and appears to be influenced by the same factors mentioned above. No definite "turning point" was observed to occur, as in the case of many animal diseases.

The above conclusions are based on studies of 12 strains of the potato blackleg parasite, including the 4 "species" originally described as being the cause of the disease.

The strains studied were observed to be morphologically similar. That they were very much alike in their cultural characteristics and physiology was abundantly proven by extensive comparative studies (see data on p. 41).

¹See Descriptive Chart, Society of American Bacteriologists.

II

Quantitative determinations of carbohydrate utilization show that *Bacillus atrosepticus* cannot hydrolyze potato starch or dextrin.

This organism, however, can utilize the saccharides, glucose, fructose, galactose, sucrose, lactose, and maltose.

The several strains (9) of *B. atrosepticus* investigated were similar in respect to their ability to hydrolyze and utilize the saccharides presented.

Bacillus coli can utilize the carbohydrates, lactose, maltose, dextrin, and potato starch, as well as glucose, fructose, and galactose. The strain investigated did not hydrolyze sucrose, nor was the amount of the original of this sugar reduced.

Bacillus vulgatus can consume the saccharides, sucrose, maltose, dextrin, and potato starch, as well as glucose, fructose, and galactose. The strain investigated could not hydrolyze nor consume lactose.

Under the conditions of the experiments a carbohydrate concentration of 0.25 per cent, 0.4 per cent, and 0.5 per cent would have furnished an ample supply for *Bacillus atrosepticus*, *B. coli*, and *B. vulgatus*, respectively.

The yeast species investigated utilized only glucose and fructose, of the carbohydrates presented. Both these saccharides were completely removed by this organism in less than 6 days, under the conditions of the experiment.

The saccharides, sucrose, lactose, maltose, dextrin, and potato starch, were not hydrolyzed appreciably if sterilized in the autoclave when dissolved in the Dunham's solution employed.

The disaccharides, sucrose, lactose, and maltose, were hydrolyzed with about equal rapidity in both the autoclave and the steamer when a very small amount of a mineral acid (0.4 per cent normal HCl) was added to "adjust" the Dunham's solution.

ACKNOWLEDGMENTS

The author wishes to express his most sincere appreciation of the cooperation and assistance accorded by many colleagues and workers. Especial thanks are due to Dr. B. M. Duggar, Dr.

- _____, _____, and Sawyer, G. C. ('16). Studies on the formation and translocation of carbohydrates in plants. *Ibid.* 6:255-384. 1916.
- Delacroix, G. ('01). Sur une maladie bacterienne de la pomme de terre. *Compt. Rend. Acad. Paris* 133: 417-419. 1901.
- _____, ('01a). Rapport sur une maladie bacterienne nouvelle de la pomme de terre. *Min. de l'Agr. Bul.* 5: 1031-1033. 1901.
- _____, ('01b). Contributions à l'étude d'une maladie nouvelle de la pomme de terre produite par le *Bacillus solanicola* nov. sp. *Compt. Rend. Acad. Paris* 133:1030-1032. 1901.
- _____, ('06). Sur une maladie de la pomme de terre produite par *Bacillus phytophthorus* (Frank) Appel. *Ibid.* 143: 383-384. 1906.
- Frank, A. B. ('97). *Kampfbuch gegen die Schädlinge unserer Feldfruchte.* 308 pp. 20 pl. 46 f. Berlin, 1897.
- _____, ('98). Untersuchungen über die verschieden en Erreger der Kartoffelfaule. *Ber. d. deut. bot. Ges.* 16:273-289. 1898.
- _____, ('99). Die Bakterienkrankheiten der Kartoffeln. *Centralbl. f. Bakt. II.* 5:98-102, 134-139. f. 1-3. 1899.
- van Hall, C. J. J. ('02). *Bijdragen tot de Kennis der Bakteriële Plantenziekten.* Inaug. Diss. Univ. Amsterdam. Coöperatieve Druckerij-Vereeniging "Plantyn" Amsterdam. 198 pp. May 12, 1902.
- _____, ('03). *Bijdragen tot de Kennis der Bakteriële Plantenziekten.* (Beiträge zur Kenntnis der Bakteriellen Pflanzenkrankheiten). *Zeitschr. f. Pflanzenkr.* 13:116-118. 1903.
- Hallier, E. ('78). *Die Parasiten der Infektionskrankheiten bei Menschen, Thieren und Pflanzen: I. Die Plastiden der niederen Pflanzen.* 92 pp. 4 pl. Leipzig, 1878.
- Harding, H. A. ('10). The constancy of certain physiological characters in the classification of bacteria. *N. Y. Agr. Exp. Sta., Tech. Bul.* 13:1-41. 1910.
- Harrison, F. C. ('07). A bacterial rot of the potato, caused by *Bacillus solanisaprus*. *Centralbl. f. Bakt. II.* 17:34-39, 120-128, 166-174, 384-395. pl. 1-3. 1907.
- Hegyi, D. ('10). Einige Beobachtungen betreffs der Schwarzbeinigkeit der Kartoffel. *Zeitschr. f. Pflanzenkr.* 20:79-81. 1910.
- International Botanical Congress of Vienna ('05). International rules of botanical nomenclature adopted by the International Botanical Congress of Vienna. 99 pp. 1905.
- International Botanical Congress of Bruxelles ('10). International rules of botanical nomenclature adopted by the International Botanical Congress of Bruxelles. 58 pp. 1910.
- Iwanoff, K. S. ('99). Ueber die Kartoffelbakteriosis in der Umgegend St. Petersburg im Jahre 1898. *Zeitschr. f. Pflanzenkr.* 9:129-131. 1899.
- Jennison, H. M. ('21). *Bacillus atrosepicius* van Hall, the cause of the blackleg disease of Irish potatoes. *Phytopath.* 11:104. 1921.
- Jensen, H. ('00). Versuche über Bakterienkrankheiten bei Kartoffeln. *Centralbl. f. Bakt. II.* 6:641-648. 1900.
- Johnson, T. ('07). Some injurious fungi found in Ireland. *Roy. Dublin Soc., Econ. Proc.* 1:345-370. pl. 32-35. f. 1-5. 1907.
- Jones, D. H. ('17). Some bacterial diseases of vegetables found in Ontario. *Ont. Dept. Agr. Bul.* 240:1-24. 1917.
- Jones, L. R. ('05). Disease resistance of potatoes. *U. S. Dept. Agr., Bur. Pl. Ind. Bul.* 87:1-39. 1905.
- _____, ('07). The blackleg disease of potatoes. *Vt. Agr. Exp. Sta., Rept.* 19:257-265. 1907.
- Kligler, I. J. ('14). Indol productions by bacteria of colon-typhoid group. *Jour. Infect. Dis.* 14:81-86. 1914.
- Kramer, E. ('91). Bakteriologische Untersuchungen über Nassfaule der Kartoffeln. *Oesterr. landwirtsch. Centralbl.* 1:11-26. 1891.
- Laurent, E. ('99). *Recherches experimentales sur les maladies des plantes.* *Inst. Pasteur, Ann.* 13:1-48. 1899.
- Lewis, H. C. ('15). On the detection and estimation of bacterial indol and observations on intercurrent phenomena. *Jour. Path. and Bact.* 19:429-443. 1915.

- Ling, A. R., and Rendle, T. ('05). The volumetric determination of reducing sugars. I. Method and discussion. *The Analyst* **30**:182-190. 1905.
- , and Jones, G. C. ('08). *Ibid.* II. The limits and accuracy of the method under standard conditions. *Ibid.* **33**:160-167. 1908.
- Morse, W. J. ('07). Potato diseases in 1907. *Me. Agr. Exp. Sta. Bul.* 149:287-330. *f.* 46-50. 1907.
- , ('09). Blackleg, a bacterial disease of Irish potato. *Ibid.* 174:309-328. 1909.
- , ('11). Control of the blackleg or blackstem disease of the potato. *Ibid.* 194:201-228. *f.* 93. 1911.
- , ('16). Potato growing and potato diseases from Maine to California. *Me. Agr. Exp. Sta., Document* 531:1-20. 1916.
- , ('17). Studies upon the blackleg disease of the potato, with special reference to the relationship of the causal organisms. *Jour. Agr. Res.* **8**:79-126. 1917.
- Murphy, P. A. ('16). The blackleg disease of potatoes. *Can. Dept. Agr., Div. Bot. Circ.* 11:1-9. *1 pl.* 1916.
- Olsen, J. C. ('10). Quantitative chemical analysis. 555 pp. 1910.
- Paine, S. G. ('17). Studies in bacteriosis I. "Blackleg" of the potato. *Jour. Agr. Sci.* **8**:480-494. 1917.
- Pethybridge, G. H., and Murphy, P. A. ('10). A bacterial disease of the potato plant in Ireland and the organism causing it. *Nature* **85**:296. 1910.
- , ('11). *Ibid.* *Proc. Roy. Irish Acad.*, **Proc.** **29B**:1-37. 1911.
- Plimmer, R. H. A. ('15). Practical organic and bio-chemistry. 636 pp. 1915.
- Prillieux, E. B. ('95-'97). *Maladies des plantes agricoles*. 2 vols. Paris, 1895-'97.
- , et Delacroix, G. ('90). La gangrene de la tige de la pomme de terre, maladie bacillaire. *Compt. Rend. Acad. Paris* **111**:208-210. Paris, 1890.
- Prunet, A. ('02). Les maladies bacteriennes de la pomme de terre. *Rev. Vit.* **17**:379-385. 1902.
- Ramsey, G. B. ('19). Viability of the potato blackleg organism. *Phytopath.* **9**:285-288. 1919.
- Reinke, J., und Berthold, G. D. W. ('79). Zersetzung der Kartoffel durch Pilze. *Untersuch. Bot. Lab. Göttingen. Heft.* 1. 100 pp. *pl.* 1-9. Berlin, 1879.
- Rosenbaum, J., and Ramsey, G. B. ('18). Influence of temperature and precipitation on the blackleg of potato. *Jour. Agr. Res.* **13**:507-513. 1918.
- Shaffer, P. A. ('14). On the determination of sugar in blood. *Jour. Biol. Chem.* **19**:285-295. 1914.
- , and Hartmann, A. F. ('21). The iodometric determination of copper and its use in sugar analysis. II. Methods for the determination of reducing sugars in blood, urine, milk and other solutions. *Ibid.* **45**:349-364; 365-390. 1921.
- Shapovalov, N., and Edson, H. A. ('19). Wound cork formation in the potato in relation to seed piece decay. *Phytopath.* **9**:483-496. *f.* 1-3. 1919.
- , ('21). Blackleg potato tuber-rot under irrigation. *Jour. Agr. Res.* **22**:81-92. *pl. A. f.* 13-16. 1921.
- Smith, E. F. ('96). Bacterial disease of tomato, eggplant and Irish potato (*Bacillus solanacearum* n. sp.). *U. S. Dept. Agr., Div. Veg. Physiol. and Path. Bul.* 12:1-26. *pl.* 1-2. 1896.
- , ('05, '11, '14). Bacteria in relation to plant diseases. *Carnegie Inst. Wash., Publ.* **27**¹⁻⁴. 1905, '11, '14.
- , ('10). *Bacillus phytophthorus* Appel. *Science N. S.* **31**:748-749. 1910.
- , ('20). An introduction to bacterial diseases of plants. 688 pp. 1920.
- Tyron, H. ('94). A new potato disease. *Queensland Dept. Agr., Ann. Rept.* **1893**:2-3. 1894.
- , ('99). Potato disease. *Queensland Agr. Jour.* **5**:57-63. 1899.
- Wehmer, C. ('98). Untersuchungen über Kartoffelkrankheiten. III. Die Bakterienfaule der Knollen (Nassfaule). *Centralbl. f. Bakt.* **II.** **4**:540-546, 570, 627-635, 694-700, 734-746, 764-770, 795-805. *pl.* 10-11. 1898.

- Winslow, C-E. A., Broadhurst, J., Buchanan, R. E., Krumwiede, C. Jr., Rogers, L. A., and Smith, G. H. ('20). The families and genera of the bacteria. Jour. Bact. 5:191-229. 1920.
- Wolf, F. A., and Foster, A. C., ('21). Studies on the physiology of some plant pathogenic bacteria. V. The fermentative activity of some plant pathogenic bacteria in relation to hydrogen ion concentration. N. C. Agr. Exp. Sta., Tech. Bul. 20:25-43. 1921.

EXPLANATION OF PLATE


PLATE 1

Fig. 1. Affected plant showing characteristic blackening of stalk and lower branches.

Fig. 2. Similarly affected plant. Lower parts of main stalk and one of the lateral branches cut away to show destruction and blackening of medullary region.

Fig. 3. Young plants prostrated by the disease. Note external signs at base of stalks. The pathogen moved upward into the stalks from the seed piece.

Fig. 4. Median longitudinal section of a recently infected stem showing disease advancing in the pith.



JENNISON—POTATO BLACKLEG

EXPLANATION OF PLATE

PLATE 2


Fig. 1. External view of tuber selected to illustrate characteristics of the rot as well as extent to which it may be manifest externally.

Fig. 2. Internal view of same. The necrosed tissues were quite dry and cheesy.

Fig. 3. External view of affected tuber. The tuber appears to be quite sound except for slight blackening and shrinking of the stolon.

Fig. 4. Internal view of same tuber. Note extent of lesion.

Fig. 5. Apparatus and materials used for determination of sugar consumption.



JENNISON—POTATO BLACKLEG

STUDIES OF SOUTH AMERICAN SENECIOS—I¹

J. M. GREENMAN

*Curator of the Herbarium of the Missouri Botanical Garden.
Professor in the Henry Shaw School of Botany
of Washington University.*

The representation of the South American flora in American herbaria has been greatly augmented during the past ten years by several scientific expeditions to South America and by material acquired from botanical correspondents and experienced collectors residing in South America.

Dr. N. L. Britton, Director of the New York Botanical Garden, has generously submitted to the writer for study the extensive suite of specimens of *Senecio*, collected in Colombia by Dr. H. H. Rusby and Dr. F. W. Pennell in 1917 and 1918; and Dr. J. N. Rose, of the United States National Museum, has placed in my hands the excellent series of specimens of this genus obtained by him and his associates on the several expeditions to South America in connection with his studies of the Cactaceae. Dr. B. L. Robinson, Curator of the Gray Herbarium of Harvard University, Mr. William R. Maxon, of the United States National Herbarium, and Dr. Charles F. Millspaugh, of the Field Museum of Natural History, have kindly loaned valuable material of this group. Dr. H. H. Rusby has given valued information and supplied material of certain specimens collected by Miguel Bang. To all these gentlemen the writer is under obligation and takes this opportunity to express his personal gratitude.

A critical study of this relatively large assemblage of specimens of South American *Senecios* has made it possible to identify many of the earlier species published by Kunth, De Candolle, Weddell, and others, as well as many of the more recent species described by Klatt, Hieronymus, and contemporary authors. It has seemed advisable to record a partial list of the species identified and to present descriptions of those species which appear not to have been previously described.

Senecio aberrans Greenm., sp. nov.

Plate 3.

Herba erecta elata; caule stricto striato juventate subarachnoideo denique plus minusve glabrato usque ad inflorescentias folioso; foliis alternis petiolatis oblongo-lanceolatis 10-15 cm. longis 2-4 cm. latis acuminatis acutis leviter sinuato-callososerratis basi rotundatis vel cuneatis juventate utrinque arach-

¹ Issued September 28, 1923.

noideo-tomentulosus supra mox glabratis reticulato-nervatis et subinde paulo bullatis, medio nervio venisque subtus prominulis; petiolis 5-7 mm. longis; inflorescentiis terminalibus multicapitatis pyramido-paniculatis, ramis paniculae subracemosis; capitulis heterogamis 12-15 mm. altis radiatis; involucris anguste campanulatis calyculatis; involucri bracteis saepe 13 lineari-lanceolatis 10-12 mm. longis acutis penicillatis glabris vel parce puberulentis, marginibus scariosis; floribus ligulatis plerumque 8, tubo gracile 7 mm. longo, ligula anguste oblonga circiter 7 mm. longa 2 mm. lata 5-nervata flava; floribus disci 12-15 tubulo-campanulatis circiter 1 cm. longis flavibus, limbo basi sensim ampliatio aequaliter 5-dentato; pappi setis circiter 1 cm. longis fuscentibus; achaeniis glabris.

Tall erect herb; stem strict, striate, subarachnoid in the early stages, later more or less glabrate, leafy to the inflorescence; leaves alternate, petiolate, oblong-lanceolate, 10-15 cm. long, 2-4 cm. broad, acuminate, acute, shallowly sinuate-callos-serrate, rotund or cuneate at the base, arachnoid-tomentulose on both surfaces in the early stages, soon glabrate, reticulately nerved, and somewhat bullate above, the median nerve and lateral veins conspicuous beneath; petioles 5-7 mm. long; inflorescence a terminal many-headed pyramidal panicle; heads heterogamous, 12-15 mm. high, radiate; involucre narrowly campanulate, calyculate; bracts of the involucre usually 13, linear-lanceolate, 10-12 mm. long, acute, penicillate, glabrous or sparingly puberulent, margins scarious; ligulate flowers usually 8, tube slender, 7 mm. long, ligule narrowly oblong, about 7 mm. long, 2 mm. broad, 5-nerved, yellow; flowers of the disk 12-15, tubular-campanulate, about 1 cm. long, yellow, the limb gradually amplified from the base, equally 5-dentate; setae of the pappus about 1 cm. long, tawny; achenes glabrous.

Colombia: depression in prairie, Mariquita, Department of Tolima, alt. 250-300 m., January 7, 1918, *Pennell 3636* (N. Y. Bot. Gard. Herb., Mo. Bot. Gard. Herb., and U. S. Nat. Herb.), TYPE; Honda, August, 1919, *Bro. Ariste-Joseph A 356* (U. S. Nat. Herb.); without definite locality, coll. of 1917, *Rusby & Pennell* (N. Y. Bot. Gard. Herb.).

This species is somewhat aberrant in the genus *Senecio* on account of the relatively long and slightly conical-tipped or obtusish style-branches. In general, however, it conforms with the technical characters of the genus.

Senecio abietinus Willd. ex Wedd. Chlor. And. 1:100. 1855-57.

Colombia: Guadalupe, near Bogotá, September, 1907, *Bro. Ariste-Joseph A28* (U. S. Nat. Herb.); Verjón, September, 1917, *Bro. Ariste-Joseph A154* (U. S. Nat. Herb. and Mo. Bot. Gard. Herb.); without definite locality or date of collection, *Bro. Ariste-Joseph* (Gray Herb., Mo. Bot. Gard. Herb., and U. S. Nat. Herb. No. 888284); bushy slope, above Bogotá, Department of Cundinamarca, alt. 2200-2800 m., August 16, 1917, *Rusby & Pennell 1289* (N. Y. Bot. Gard. Herb. and Field Mus. Herb.); open rocky mountain slope, Chapinero, near Bogotá, Department of Cundinamarca, alt. 3000-3100 m., September 18-23, 1917, *Pennell 2019* (N. Y. Bot. Gard. Herb., Mo. Bot. Gard. Herb., and U. S. Nat. Herb.); Páramo de Choachi, near Bogotá, alt. 3700 m., August 8, 1922, *Killip & Bro. Ariste-Joseph 11928* (U. S. Nat. Herb.); Cundinamarca, coll. of 1883, *Lehmann 2406* (Gray Herb.); coll. of 1861, ex Herb. *J. de Parsaval-Grandmaison 92* (Gray Herb.).

Venezuela: Páramo de Tama, head of Tachira River, alt. 2440 m., March, 1911, *Wilfred H. Osgood* (Field Mus. Nat. Hist. Herb.).

Senecio adenophyllus Meyen & Walp. in Nov. Act. Nat. Cur. 19: Suppl. 1, p. 282. 1843; Walp. Rep. 6:271. 1846-47; Wedd. Chlor. And. 1:112. 1855-57.

Bolivia: vicinity of Oruro, August 18, 1914, *Mr. & Mrs. J. N. Rose 18932* (U. S. Nat. Herb. and Mo. Bot. Gard. Herb.).

Peru: below Pampa del Arrieros, August 23, 1914, *Mr. & Mrs. J. N. Rose 18951* (U. S. Nat. Herb. and Mo. Bot. Gard. Herb.).

The leaves of this species are apparently quite variable in size and outline. Walper's description reads "foliis sessilibus confertis ovato-lanceolatis acuminatis grosse serratis vel subintegris lineari-lanceolatis." In the Gray Herbarium there is a drawing of Meyen's original material from Peru which shows a narrowly lanceolate, coarsely and irregularly dentate form of leaf. Although the leaves on the specimens above cited are narrower than they would appear to be in the type, yet the specimens collected by Mr. and Mrs. Rose in all probability represent a narrow-leaved form of the above species.

Senecio adenotrichus DC. Prodr. 6:416. 1837.

Chile: Quillota, *Bertero 1044* (Mo. Bot. Gard. Herb.); without locality, *Styles* (Phil. Acad. Nat. Sci. Herb.); Palos Quernados, October 4, 1914, *Mr. & Mrs. J. N. Rose 19187* (U. S. Nat. Herb.).

and Mo. Bot. Gard. Herb.); vicinity of Choapa, and vicinity of Illapel, October 6, 1914, *Mr. & Mrs. J. N. Rose 19193, 19362, 19237* (U. S. Nat. Herb.); San Felipe, September 25, 1919, *Holway & Holway 67* (Gray Herb.).

Senecio alternifolius Greenm., comb. nov.

Gynoxis alternifolia Schz. Bip. in Bull. Soc. Bot. Fr. 12:80. 1865, name only; in Linnaea 34:531. 1866, name only; Rusby in Mem. Torr. Bot. Club 6:67. 1896, description.

Bolivia: vicinity of Sorata, alt. 3000-3200 m., May, 1859, *Mandon 131* (Gray Herb.); vicinity of Mapiri, alt. 2435 m., September, 1892, *Bang 1574* (Mo. Bot. Gard. Herb.); Unduavi, September, 1894, *Bang 2477* (Mo. Bot. Gard. Herb. and Gray Herb.), distributed as "*Eupatorium trichotomum* Sch. Bip."; Unduavi, Noryungas, alt. 3300 m., November, 1910, *Buchtien 3078* (N. Y. Bot. Gard. Herb.).

The two morphological characters, namely, opposite leaves and conical tipped style-branches, ordinarily relied upon to distinguish *Gynoxis* from *Senecio*, are at best very unsatisfactory. The latter character does not hold, since the same type of style-branch occurs in a number of species which are accepted unqualifiedly as true *Senecios*. The one character separating these two genera is the leaf arrangement. In only one case, among the several collections cited above, namely, *Bang's* No. 1574, do the leaves approach an opposite position on the stem, and even here they are only subopposite. In all other regards the species in question is a typical *Senecio*, hence it is transferred to that genus.

Senecio andicola Turcz. Bull. Soc. Nat. Mosc. 24²:91. 1851.

S. vernicosus Schz. Bip. var. *major* Wedd. Chlor. And. 1:94. 1855-1857.

S. ledifolius Schz. Bip. acc. to Wedd. l. c.

Colombia: moist open places, Páramo de Ruiz, Department of Tolima, alt. 3400-3700 m., December 16-17, 1917, *Pennell 3076* (N. Y. Bot. Gard. Herb., U. S. Nat. Herb., and Mo. Bot. Gard. Herb.).

A study of additional material may show that this species should be merged with *S. ledifolius* (HBK.) DC., but at present it seems desirable to regard it as specifically distinct, on account of the larger leaves and broader calyculate bracteoles of the involucre.

Senecio apiculatus Schz. Bip. ex Wedd. Chlor. And. 1:128. 1855-57.

Venezuela: Páramo de Timotes, State of Táchira, alt. 3000-3500 m., March, 1910, *Jahn 6* (U. S. Nat. Herb.); Páramo Piedras Blancas, alt. 4000 m., March 27, 1915, *Jahn 426* (U. S. Nat. Herb.); Páramo de La Sal, Mérida, alt. 3400 m., September, 1921, *Jahn 556* (U. S. Nat. Herb.); Páramo de Timotes, Mérida, alt. 3600 m., January 22, 1922, *Jahn 789* (U. S. Nat. Herb.).

This species is known in Venezuela under the common name of "Romerito cenizo."

Senecio arboreus Greenm., comb. nov.

Cacalia arborea HBK. Nov. Gen. & Sp. 4:163, pl. 359. 1820.

Colombia: shrub zone, below Páramo del Chaquiro, Cordillera Occidental, Department of Bolívar, alt. 2800-3100 m., February 24, 1918, *Pennell 4313* (N. Y. Bot. Gard. Herb.).

Senecio aridus Greenm., sp. nov.

Plate 4.

Herbaceous perennis ubique floccoso-tomentosus; caule erecto vel suberecto circiter 3 dm. alto plus minusve glabrato; foliis alternis inferioribus oblanceolatis 3.5-8 cm. longis 4-10 mm. latis acutis vel obtusis calloso-denticulatis vel integris marginibus revolutis juventute utrinque floccoso-tomentosis supra plus minusve glabratis, eis caulinis lanceolatis sessilibus et brevi-decurrentibus; inflorescentiis terminalibus paucicapitatis magnis radiatis; involucri campanulatis calyculatis; involucri bracteae 21 lineari-lanceolatis circiter 13 mm. longis 2-3 mm. latis parce tomentulosae ad apicem penicillatis; floribus liguliferis 15-20, ligulis late oblongis 13-15 mm. longis 5-7 mm. latis pallide flavibus; floribus disci circiter 125; achaeniis glabris.

Herbaceous perennial, floccose-tomentulose throughout; stem erect or suberect, about 3 dm. high, more or less glabrate; leaves alternate, the lowermost oblanceolate, 3.5-8 cm. long, 4-10 mm. broad, acute or obtuse, callous-denticulate, or entire, revolute-margined, in the early stages floccose-tomentose on both surfaces, more or less glabrate above; upper stem-leaves lanceolate, sessile and short-decurrent; inflorescence terminal, few-headed; heads large, radiate; involucre campanulate, calyculate; bracts of involucre 21, linear-lanceolate, about 13 mm. long, 2-3 mm. broad, sparingly tomentulose, penicillate at the apex; ray-flowers 15-20, rays broadly oblong, 13-15 mm. long, 5-7 mm. broad, light yellow; disk-flowers about 125; achenes glabrous.

Colombia: dry open places, Páramo de Ruiz, Department of Tolima, alt. 3800-4300 m., December 16-17, 1917, *Pennell 3037* (N. Y. Bot. Gard. Herb. and Mo. Bot. Gard. Herb.), TYPE.

This species is related to *S. latiflorus* Wedd., but differs in being floccose-tomentulose throughout, in having much smaller and shorter leaves, and in having smaller heads with fewer flowers.

Senecio auritus Wawra Itin. Princ. S. Coburg. 2:47. 1888.

Brazil: vicinity of Itatiaya, July 26-30, 1915, *Rose & Russell* 20559 (U. S. Nat. Herb., fragment and photograph in Mo. Bot. Gard. Herb.).

Senecio brachycodon Baker in Mart. Fl. Bras. 6^a:319. 1884.

Brazil: vicinity of Itatiaya, July 26-30, 1915, *Rose & Russell* 20506 (U. S. Nat. Herb., fragment and photograph in Mo. Bot. Gard. Herb.).

Senecio Buchtienii Greenm., sp. nov.

Frutex 2-3 m. altus; ramis ramulisque striato-angulatis persistenter papilloso-hispidis; foliis alternis petiolatis lanceolatis vel lanceolato-oblongis 5-8 cm. longis 7-20 mm. latis sinuato-dentatis acutis basi cuneatis juventate parce pubescentibus denique supra glabris et atro-viridibus subtus pallidioribus nisi in nervo medio glabris; petiolis usque ad 1.5 cm. longis; inflorescentiis terminalibus vel lateralibus laxe corymboso-cymosis bracteatis fusco-pubescentibus, bracteis lanceolatis vel linearibus; capitulis heterogamis 1-1.5 cm. altis; involucris anguste campanulatis calyculatis, squamellis calyculatis linearibus 3 mm. vel minus longis; involucri bracteis 8 lineari-lanceolatis 8-9 mm. longis glabris; floribus liguliferis 5, ligulis parvis linearibus 5-6 mm. longis plerumque bidentatis flavibus; floribus disci 10-12; pappi setis albidis ad corollam subaequantibus; achaeniis glabris.

Shrub 2-3 m. high; branches and branchlets striate-angled, persistently papillose-hispid; leaves alternate, petiolate, lanceolate or lanceolate-oblong, 5-8 cm. long, 7-20 mm. broad, sinuate-dentate, acute, cuneate at the base, in the young stages sparingly pubescent, later becoming glabrous and dark green above, paler beneath and glabrous or hirsute pubescent on the midrib; petioles 1.5 cm. or less in length; inflorescence terminating the stem and lateral branches in bracteate tawny pubescent loose corymbose cymes; bracts of the inflorescence lanceolate to linear; heads heterogamous, 1-1.5 cm. high; involucre narrowly campanulate, calyculate, bracteoles linear, 3 mm. or less in length; bracts of the involucre 8, linear-lanceolate, 8-9 mm. long, glabrous; ligulate flowers 5, ligules small, linear, 5-6 mm. long,

usually bidentate, yellow; flowers of the disk 10-12; pappus white, about as long as the corolla; achenes glabrous.

Bolivia: Unduavi, Noryungas, alt. 3300 m., November, 1910, *O. Buchtien 3087* (U. S. Nat. Herb., N. Y. Bot. Gard. Herb., fragment and photograph in Mo. Bot. Gard. Herb.), TYPE.

This species appears to be most nearly related to *S. Sepium* Schz. Bip. ex Rusby in Bull. N. Y. Bot. Gard. 4:394. 1907, but differs in having lanceolate leaves, which are cuneate at the base, instead of ovate-cordate leaves, and in having persistently papillose-hispid stems.

***Senecio chaquiroensis* Greenm., sp. nov.**

Frutex scandens; ramis ramulisque teretibus dense fulvo-hispidis; foliis alternis petiolatis elliptico-oblongis 2.5-5 cm. longis 1.2-2 cm. latis mucronatis vel obtusis callosio-denticulatis vel subintegris supra glabris subtus pallidioribus et valde reticulato-venosis plus minusve puberulentis, marginibus revolutis; petiolis 5-7 mm. longis canaliculatis puberulentis vel glabris; inflorescentiis paniculatis fulvo-hispidis; capitulis homogamis; involucri brevi-calyculatis anguste campanulatis, bracteolis calyculatis triangulari-lanceolatis 1-2 mm. longis acutis ciliatis; bracteis involucri 8 anguste oblongis vel lineari-lanceolatis circiter 5 mm. longis extrinsecus glabris acutis penicillatisque; floribus 12; corollis 3-4 mm. longis glabris luteo-albidis pappi setis albidis subaequantibus; achaeniis glabris.

Stem scandent, ligneous; branches and branchlets terete, densely tawny hispid; leaves alternate, petiolate, elliptic-oblong, 2.5-5 cm. long, 1.2-2 cm. broad, mucronate or obtuse, callous-denticulate or subentire, glabrous above, paler and strongly reticulately veined and more or less puberulent beneath, margins revolute; petioles 5-7 mm. long, canaliculate, puberulent or glabrous; inflorescence paniculate, tawny hispid; heads homogamous; involucre short-calyculate, narrowly campanulate, the calyculate bracteoles triangular-lanceolate, 1-2 mm. long, acute, ciliate; bracts of the involucre 8, narrowly oblong or linear-lanceolate, about 5 mm. long, glabrous on the outer surface, acute and penicillate; flowers 12; corollas 3-4 mm. long, glabrous, about as long as the white setae of the pappus; achenes glabrous.

Colombia: shrub zone, below Páramo de Chaquiro, Cordillera Occidental, Department of Bolívar, alt. 2800-3100 m., February 24, 1918, *Pennell 4290* (N. Y. Bot. Gard. Herb. and Mo. Bot. Gard. Herb.).

This species is related to *S. theaeifolius* Benth., but differs in having the branches and inflorescence distinctly tawny and densely hispid pubescent, in having the lower leaf-surface more or less puberulent, especially on the midrib, lateral nerves and reticulated veins, and in the somewhat shorter involucre and smaller flowers.

Senecio Chinogeton Wedd. forma **macrocephalus** Hieron. in Engl. Bot. Jahrb. 19:64. 1894.

Colombia: in bushy mountain valley, Rio San Cristobal, near Bogotá, Department of Cundinamarca, alt. 3000-3300 m., September 20-26, 1917, *Pennell 2047* (N. Y. Bot. Gard. Herb., Mo. Bot. Gard. Herb., and U. S. Nat. Herb.).

The specimens under Pennell's No. 2047 show only the complete inflorescence with mature, discoid, nodding heads, and a few of the uppermost stem-leaves. From the original description of the above species they would seem to be conspecific and probably represent the form "*macrocephala*", briefly characterized by Professor Hieronymus.

Senecio clavifolius Rusby in Mem. Torr. Bot. Club 3:64. 1893.

S. attenuatus var. *microphyllus* Britt. in Bull. Torr. Bot. Club 19:264. 1892.

Bolivia: Talca Chugiaguillo, April, 1890, *Bang 792* (U. S. Nat. Herb., Phil. Acad. Nat. Sci. Herb., Field Mus. Nat. Hist. Herb., and Mo. Bot. Gard. Herb.); near La Paz, alt. 3500 m., October, 1885, *Rusby 1691* (U. S. Nat. Herb., Phil. Acad. Nat. Sci. Herb., and Field Mus. Nat. Hist. Herb.).

Senecio comosus Schz. Bip. in Bonplandia 4:52, 55. 1856, nomen subnudum; Bull. Soc. Bot. Fr. 12:180. 1865, name only; Linnaea 34:531. 1866, name only; Wedd. Chlor. And. 1:129. 1855-57, description.

Bolivia: coll. of 1856-61, *Mandon 136* (Gray Herb.); vicinity of Cochabamba, coll. of 1891, *Bang 1240* (Mo. Bot. Gard. Herb., Phil. Acad. Nat. Sci. Herb., and Gray Herb.); Cochabamba, April 13-21, 1892, *Otto Kuntze* (U. S. Nat. Herb., No. 702117).

Mr. Bang's No. 1240 was taken to represent *Senecio culcitoides* Schz. Bip., which it resembles habitally, and distributed as that species. The specific name, however, was inadvertently written on the label as "*calcitoides*," and the plant was referred to in Mem. Torr. Bot. Club 6:65. 1896 as "*Senecio culcitroides* Wedd." The heads of the Bang plant are distinctly radiate, not discoid as in *L. culcitoides* Schz. Bip. The discoid heads and the very broad, conspicuous, almost glabrous midvein on the lower

side of the leaf of *S. culcitoides* serve to separate it from *S. comosus*.

Senecio culcitoides Schz. Bip. in Bonplandia 4:55. 1856, name only and by error as "*cultitoides*"; in Lechler, Berb. Am. Austr. 57. 1857, name only; in Wedd. Chlor. And. 1:103. 1855-57, description; in Bull. Soc. Bot. Fr. 12:80. 1865, name only; in Linnaea 34:530. 1866, name only.

Colombia: dry open places, Páramo de Ruiz, Department of Tolima, alt. 3500-4100 m., December 16-17, 1917, Pennell 3026 (N. Y. Bot. Gard. Herb. and Mo. Bot. Gard. Herb.).

Bolivia: Mandon 116 (Gray Herb.).

Dr. Pennell's specimen here cited agrees in all essential details with the original description of the above species except in the number of involueral bracts which, according to Weddell, is 12-15. In the specimen at hand the involucre consists definitely of 21 bracts, disposed in a single series. The subtending bracteoles are relatively conspicuous, yet they do not form a part of the true uniseriate involucre; and on this account the plant in question cannot be referred properly to *Culcitium*. In habit and foliar characters Dr. Pennell's specimen simulates *Culcitium longifolium* Turcz., but that species is described as having a multiseriate involucre.

Senecio decompositus Schz. Bip. ex Hieron. in Engl. Bot. Jahrb. 28:634. 1901.

Colombia: Santa Marta Expedition, 1898-9, Herbert H. Smith 1989 (Mo. Bot. Gard. Herb., N. Y. Bot. Gard. Herb., and Field Mus. Nat. Hist. Herb.).

The specimens here cited agree in all details with the original description of the above species by Professor Hieronymus. The rather leafy panicle in the specimens at hand varies from 2 to 3 dm. in length and from 1 to 2.5 dm. in breadth. The heads are very numerous, relatively small, and bear regularly, as far as examined, 10 flowers. The collector's comments on the specimens collected read,—"A loosely twining vine, to 20 feet. Moderately common on the Sierra del Libano thickets or dry forest on ridges, 6000-7000 feet. Begins to flower about Jan. 15. The specimens were collected Jan. 25, at 6000 feet. Flowers rich yellow; leaves polished. This is one of the most showy species of the order; a plant in full bloom is a fine sight."

Senecio ericaefolius Benth. Pl. Hartw. 208. 1845.

Peru: western Cordillera, opposite Huancabamba, alt. 2400-

2900 m., September 26, 1911, *C. H. T. Townsend A207* (Field Mus. Nat. Hist. Herb.).

Senecio formosus HBK. Nov. Gen. & Sp. 4:177. 1820; DC. Prodr. 6:428. 1837; Wedd. Chlor. And. 1:125. 1855-57, excl. *γ. subtruncatus*.

S. Tabacon Turcz. in Bull. Soc. Nat. Mosc. 24²:91. 1851.

Colombia: *Moritz 1400* (Gray Herb.); Páramo de Buena Vista, Huila group, Central Cordillera, State of Cauca, alt. 3000-3600 m., January, 1906, *Jahn 1117* (U. S. Nat. Herb.); moist bushy valley, Páramo de Choachi, near Bogotá, Department of Cundinamarca, alt. 3000-3100 m., September 27, 1917, *Pennell 2237* (N. Y. Bot. Gard. Herb.); in swale, "Rosalito", near Páramo de Ruiz, Department of Tolima, alt. 2800-3100 m., December 15-17, 1917, *Pennell 2943* (N. Y. Bot. Gard. Herb. and Mo. Bot. Gard. Herb.); Laguna de Buitrago, alt. 3000 m., December, 1919, *Bro. Ariste-Joseph A 503, A 504, A 505* (U. S. Nat. Herb.); Laguna Coche, May 3, 1876, *Andre 3092* (N. Y. Bot. Gard. Herb.).

Venezuela: Páramo de la Cristalina, State of Trujillo, alt. 2900 m., December 20, 1910, *Jahn 13* (U. S. Nat. Herb.).

When *Senecio formosus* HBK. was published originally it was doubtfully attributed to "Nova Hispania," or Mexico. Weddell pointed out that the species is a South American one, and that its erroneous assignment to Mexico was probably due to a confusion of labels accompanying the original material. Weddell's treatment of the species, however, would indicate that he regarded it as a polymorphic species, since he included *S. Tabacon* Turcz. as a synonym and then characterized three varieties, namely, *angustifolius*, *latifolius*, and *subtruncatus*. To the writer the last-named variety seems to merit specific rank and is so treated in this paper. The specimens here cited as representing *S. formosus* show considerable variation in leaf-outline and in the degree of pubescence on the achenes. In Pennell's No. 2943 the achenes are but slightly hirtellous, while in all the others the hirtellous character is very distinct. The species is well marked by the glandular-villous hairs on stem, leaves, and involucre, and by the purple rays and numerous disk-flowers.

Senecio genuflexus Greenm., sp. nov.

Frutex scandens; caulibus vel ramis gracilibus glabris plus minusve flexuosis vel geniculatis; foliis alternis petiolatis oblongo-lanceolatis vel ovato-lanceolatis 2.5-6 cm. longis 1-2.5 cm. latis acutis vel acuminatis submucronatisque integris vel subin-

tegris basi rotundatis vel cuneatis supra glabris subtus juventate parce arachnoideo-puberulentis mox glabrat; petiolis .5–2 cm. longis paululo alatis et basi abrupte ampliatis auriculato-amplexicaulibusque; inflorescentiis terminalibus multicapitatis paniculatis; rhachis paniculatae geniculatis glabris vel parcissime puberulentis; capitulis discoideis 6–8 mm. altis; involucris anguste campanulatis calyculatis; bracteis involucri saepe 8 linearilanceolatis 3–3.5 mm. longis glabris; floribus disci 12–14; pappi setis albidis; acheniis glabris.

Scandent shrub; stems slender, glabrous, more or less flexuous or geniculate; leaves alternate, petiolate, oblong-lanceolate to ovate-lanceolate, 2.5–6 cm. long, 1–2.5 cm. broad, acute or acuminate and submucronate, entire or subentire, rotund or cuneate at the base, glabrous above, in the early stages very sparingly arachnoid-puberulent beneath but soon glabrate; petioles .5–2 cm. long, narrowly winged, abruptly expanded at the base and auriculate-amplexicaul; inflorescence a terminal many-headed panicle; rhachis of the panicle geniculate, glabrous or very sparingly puberulent; heads discoid, 6–8 mm. high; involucre narrowly campanulate, calyculate; bracts of the involucre usually 8, linear-lanceolate, 3–3.5 mm. long, acute, glabrous; flowers of the disk 12–14; bristles of the pappus white; achenes glabrous.

Colombia: thickets on ridges, Sierra del Libano, Santa Marta, alt. 1800–2000 m., January 19, 1899, *Herbert H. Smith* 1987 (Mo. Bot. Gard. Herb., TYPE; Gray Herb., in part; U. S. Nat. Herb., in part; N. Y. Bot. Gard. Herb., in part; Field Mus. Nat. Hist. Herb.).

The plant here cited was distributed under the name "*Senecio decompositus* Schz. Bip." with which species it was doubtfully identified, but from which it is readily distinguished by the stipule-like amplexicaul appendages at the base of the petiole and by the strongly geniculate main axis of the inflorescence.

Senecio involucratus (HBK.) DC. Prodr. 6:422. 1837; Wedd. Chlor. And. 1:93. 1855–57.

Cacalia involucrata HBK. Nov. Gen. & Sp. 4:166. 1820.

Aetheolaena involucrata Cass. Dict. Sci. Nat. 48:453. 1827.

Ecuador: Nabón, September 25, 1918, *Rose, Pachano & Rose* 22995 (U. S. Nat. Herb. and N. Y. Bot. Gard. Herb.).

Senecio iodopappus Schz. Bip. in Bonplandia 4:55. 1856, name only (*jodopappus*); Wedd. Chlor. And. 1:116, pl. 20, fig. B. 1855–57.

Bolivia: Lake Titicaca, May, 1910, *Buchtien 3106* (U. S. Nat. Herb.); vicinity of Oruro, August 18, 1914, *Mr. & Mrs. J. N. Rose 18926* (U. S. Nat. Herb.).

Senecio laciniatus HBK. Nov. Gen. & Sp. 4:175. 1820.

S. pimpinellifolius var. *laciniata* Hieron. in Engl. Bot. Jahrb. 28:634. 1901.

Ecuador: vicinity of Zaragura, September 28, 1918, *Rose, Pachano & Rose 23143* (U. S. Nat. Herb.).

Peru: *Mathews 87* (Gray Herb.).

Senecio latiflorus Wedd. Chlor. And. 1:125. 1855-57.

Colombia: dry, open, rocky places, Páramo de Ruiz, Department of Tolima, alt. 3700-4200 m., December 16, 17, 1917, *Pennell 3033* (N. Y. Bot. Gard. Herb., U. S. Nat. Herb., Mo. Bot. Gard. Herb., and Gray Herb.); in moist lands, Ruiz, alt. 3000-3500 m., coll. of 1918, *Dawe 749*, and *763* (N. Y. Bot. Gard. Herb.); *Triana 98* (Gray Herb.).

Senecio lophophilus Greenm., sp. nov.

Plate 5.

Fruticulus .5-1 m. altus plus minusve diffusus ubique glabrus; caulibus teretibus striatis brunneis; foliis alternis petiolatis lanceolatis 3-6 cm. longis 1-2 cm. latis acutis sinuato-dentatis basi cuneatis utrinque glabris luteo-viridibus; petiolis usque ad 5 mm. longis; inflorescentiis terminalibus dense paniculatis 7 cm. longis 5 cm. latis; capitulis brevi-pedunculatis parvis 6-7 mm. altis discoideis; involucri anguste campanulatis parce calyculatis; bracteis involucri 8 lineari-lanceolatis circiter 5 mm. longis acutis glabris stramineis; floribus 10-12, corollis 5-6 mm. longis flavibus; achaeniis glabris.

Small shrub, .5-1 m. high, more or less diffuse, glabrous throughout; stems terete, striate, brownish; leaves alternate, petiolate, lanceolate, 3-6 cm. long, 1-2 cm. broad, acute, sinuate-dentate, cuneate at the base, glabrous on both surfaces, yellowish green; petioles 5 mm. or less in length; inflorescence a dense terminal panicle, 7 cm. long, 5 cm. broad; heads short-pedunculate, small, 6-7 mm. high, discoid; involucre narrowly campanulate, sparingly calyculate; bracts of the involucre 8, linear-lanceolate, about 5 mm. long, acute, glabrous, stramineous; flowers, 10-12, corollas 5-6 mm. long, yellow; achenes glabrous.

Colombia: open lands on San Lorenzo Ridge, Santa Marta, alt. 2000-2300 m., January 27, 1899, *Herbert H. Smith 1988* (N. Y. Bot. Gard. Herb., fragment and photograph in Mo. Bot. Gard. Herb.).

In general the species here described stands near *S. prunifolius* Wedd., but it differs in leaf-outline and in being glabrous throughout.

Senecio Magnusii Hieron. in Engl. Bot. Jahrb. 28:642. 1901.

Colombia: edge of tree "island" in páramo valley, Mt. Chuscal, west of Zipaquira, Department of Cundinamarca, alt. 3200–3300 m., October 22, 1917, *Pennell 2589* (N. Y. Bot. Gard. Herb. and Mo. Bot. Gard. Herb.); forest on mountain, south of Sibate, Department of Cundinamarca, alt. 2800–3000 m., October 13–15, 1917, *Pennell 2481* (N. Y. Bot. Gard. Herb. and Mo. Bot. Gard. Herb.).

The specimens here cited agree in all essential details with the original description of *S. Magnusii* Hieron. and are hereto referred pending a comparison with type material of this species. The specimens at hand also possess several characters in common with *L. disciformis* Hieron.

Senecio medullosus Schz. Bip. in Bull. Soc. Bot. Fr. 12:80. 1865, name only; in *Linnaea* 34:531. 1866, name only.

Verisimiliter frutex erectus; caulibus ramisque teretibus vel plus minusve striato-angulatis glabris; foliis alternis petiolatis lanceolatis vel lanceolato-oblongis 5–18 cm. longis 1–4.5 cm. latis acuminatis acutis remote calloso-denticulatis vel sinuato-subseratis basi cuneatis juventate parcissime puberulentis nisi in nervo supra medio utrinque mox glabratis; petiolis circiter 1 cm. longis; inflorescentiis terminalibus paniculatis patentibus parce puberulentis; capitulis heterogamis 10–15 mm. altis; involucris campanulatis, brevi-calyculatis; bracteis involucris 13, linearilanceolatis vel lanceolato-oblongis 7–8 mm. longis 1.5–2 mm. latis acutis vel obtusis glabris viridibus vel plus minusve purpurascenscentibus; floribus liguliferis 6–8 purpurascenscentibus, tubo gracile 5–10 mm. longo curvato, ligulis parvulis oblongis circiter 5 mm. longis; floribus disci 18–20, corollis tubulo-campanulatis aequaliter 5-dentatis plus minusve purpurascenscentibus; pappi setis albidis corollo brevioribus; achaeniis striatis glabris.

Shrub, probably erect; stem and branches terete or more or less striate-angled, glabrous; leaves alternate, petiolate, lanceolate or lanceolate-oblong, 5–18 cm. long, 1–4.5 cm. broad, acuminate, acute, remotely callous-denticulate or sinuate-subserate, cuneate at the base, glabrous or very sparingly puberulent in the young stages, soon becoming glabrous on both surfaces except along the midrib above; petioles about 1 cm. long; inflorescence a terminal spreading sparingly puberulent panicle; heads heter-

ogamous, 10–15 mm. high; involucre campanulate, short-calyculate; bracts of the involucre 13, linear-lanceolate or lanceolate-oblong, 7–8 mm. long, 1.5–2 mm. broad, acute or obtuse, glabrous except at the penicillate tip, green or more or less purple; ligulate flowers 6–8, purple, tube slender, 5–10 mm. long, somewhat outwardly curved, ligules small, oblong, about 5 mm. long; disk-flowers 18–20, corollas tubular-campanulate, equally 5-dentate, more or less purple; pappus bristles white, shorter than the corollas; achenes striate, glabrous.

Bolivia: vicinity of Sorata, alt. 2900 m., October, 1858, *Mendon 147* (Gray Herb., fragment and photograph in Mo. Bot. Gard. Herb.), TYPE; Unduavi, Noryungas, alt. 3300 m., November, 1910, *Buchtien 3086* (N. Y. Bot. Gard. Herb., fragment and photograph in Mo. Bot. Gard. Herb.); San Felipe, Provincia de Sur Yungas, March 20, 1920, *Holway & Holway 627* (U. S. Nat. Herb.).

The species here described bears a superficial resemblance to *S. biacuminatus* Rusby, but differs in having purple flowers and in having a long slender tube and very small rays to the ray-flowers. It is also related apparently to *S. reflexus* HBK., which is said to have very entire reflexed leaves.

Senecio Microchaete (Benth.) Wedd. Chlor. And. 1:100. 1855-57.

Microchaete corymbosa Benth. Pl. Hartw. 196. 1845.

Colombia: near Bogotá, *Hartweg 1086* (Gray Herb.); wet páramo valley, Mt. Chuscal, west of Zipaquira, Department of Cundinamarca, alt. 3100–3300 m., October 22, 1917, *Pennell 2580* (N. Y. Bot. Gard. Herb., Field Mus. Nat. Hist. Herb., U. S. Nat. Herb., and Mo. Bot. Gard. Herb.); Guadalupe, near Bogotá, July 10, 1917, *Bro. Ariste-Joseph A14* (U. S. Nat. Herb.).

Doctor Pennell's specimens agree well with Weddell's characterization of the above species and also with the original description and specimen of *Microchaete corymbosa* Benth. on which *Senecio Microchaete* Wedd. was founded. The material also accords in all essential details with the description of *Senecio pungens* (HBK.) DC., from which it is doubtful if *S. Microchaete* Wedd. can be separated satisfactorily. It seems desirable, however, to retain Weddell's name until a careful comparison can be made with the type or authentic specimens of *L. pungens* (HBK.) DC.

Senecio modestus Wedd. Chlor. And. 1:105, pl. 18B. 1855-1857, not Phil. in Linnaea 28:745. 1856.

Bolivia: without locality, *Bang* 1890 (Mo. Bot. Gard. Herb., Phil. Acad. Nat. Sci. Herb., Gray Herb., and U. S. Nat. Herb.).

The specimen here cited agrees in all essential characters with the original description and illustration of the above species. Mr. Bang's No. 1890 was referred doubtfully by Doctor Rusby, Bull. N. Y. Bot. Gard. 4:397. 1907, to *S. rhizomatus* Rusby from which it is amply distinct.

Senecio nubigenus HBK. Nov. Gen. & Sp. 4:174. 1820.

S. pimpinellifolius var. *nubigena* (HBK.) Hieron. in Engl. Bot. Jahrb. 28:634. 1901.

Ecuador: Pichinca, alt. 3350 m., March, 1864, *Jameson* (U. S. Nat. Herb.); vicinity of Santa Rosa de Cañar, September 14, 15, 1918, *Rose & Rose* 22669 (U. S. Nat. Herb. and Gray Herb.).

The specimens here cited agree in all essential characters with the original description of the above species, although in some respects they correspond equally well to *S. pimpinellaefolius* HBK. In foliar characters, however, the material at hand seems to accord more closely with *S. nubigenus*. Further collections, upon comparison with the types, may prove the two species, as characterized by Kunth, to be conspecific.

Senecio organensis Casar. Nov. Stirp. Bras. Dec. 77; Gard. in Hook. Lond. Jour. Bot. 4:127. 1845; Baker in Mart. Fl. Bras. 6³ 320. 1884.

? *Cacalia dichroa* Bong. Mem. Imp. Acad. Sci. St. Petersburg. 1: 40, t. 7. 1831.

Brazil: vicinity of Itatiaya, July 26-30, 1915, *Rose & Russell* 20565 (U. S. Nat. Herb., fragment and photograph in Mo. Bot. Gard. Herb.).

Senecio otophorus Wedd. Chlor. And. 1:98. 1855-57.

Colombia: Páramo de Buena Vista, Huila group, Central Cordillera, State of Cauca, alt. 3000-3600 m., January, 1906, *Pittier* 1129 (U. S. Nat. Herb.); dry, open places. Páramo de Ruiz, Department of Tolima, alt. 3500-3700 m., December 16-17, 1917, *Pennell* 3005 (N. Y. Bot. Gard. Herb. and Mo. Bot. Gard. Herb.); Laguna de Buitrago, alt. 3100 m., December, 1919, *Bro. Ariste-Joseph* A 500 (U. S. Nat. Herb.).

Ecuador: Páramo de Azuay, *Spruce* (Gray Herb.); vicinity of Cañar, September 16, 1918, *Rose & Rose* 22758 (U. S. Nat. Herb.).

Senecio patens (HBK.) DC. Prodr. 6:423. 1837.

Cacalia patens HBK. Nov. Gen. & Sp. 4:164. 1820.

Colombia: in swale, "Rosalito," near Páramo de Ruiz, Department of Tolima, alt. 2800–3100 m., December 15–17, 1917, *Pennell 2971* (N. Y. Bot. Gard. Herb., U. S. Nat. Herb., and Mo. Bot. Gard. Herb.).

Senecio pellucidinervis Schz. Bip. ex Baker in Mart. Fl. Bras. 6:319. 1884.

Brazil: vicinity of Itatiaya, July 26–30, 1915, *Rose & Russell 20462, 20571* (U. S. Nat. Herb., fragment and photograph in Mo. Bot. Gard. Herb.).

Senecio Pennellii Greenm., sp. nov.

Plate 6.

Frutex scandens; caulibus teretibus striatis juventate arachnoideis mox glabratiss, brunneis; foliis alternis petiolatis ovatis vel oblongo-ovatis 10–18 cm. longis 4–8 cm. latis acutis callosodenticulatis basi leviter cordatis vel rotundatis utrinque glabris vel juventate subtus parcissime arachnoideis mox glabratiss, nervis utrinque prominenter reticulatis; petiolis 1.5–3 cm. longis; inflorescentiis terminalibus paniculatis pleuricapitatis; capitulis 12–15 mm. altis heterogamis; involucri campanulatis calyculatis; involucri bracteis 8 anguste oblongis vel oblongo-ovatis 5–6 mm. longis 1.5–3.5 mm. latis ad apicem obtusis vel subrotundatis et pubescentibus cetero glabris; floribus liguliferis 8, tubo gracile 3–3.5 mm. longo, limbo plerumque radiato vel nonnumquam irregulariter dentato flavo, floribus disci 25–30, corollis tubulo-campanulatis, flavibus; pappi setis copiosis albidis; achaeniis glabris.

Scandent shrub; stems terete, striate, arachnoid in the young stages, soon glabrate, brownish; leaves alternate, petiolate, ovate to oblong-ovate, 10–18 cm. long, 4–8 cm. broad, acute, callous-denticulate, shallowly cordate to rotund at the base, glabrous on both sides or in the young stages sparingly arachnoid beneath but soon glabrate, prominently reticulate-veined on both surfaces; petioles 1.5–3 cm. long; inflorescence a terminal many-headed panicle; heads 12–15 mm. high, heterogamous; involucre campanulate, calyculate; bracts of the involucre 8, narrowly oblong or oblong-ovate, 5–6 mm. long, 1.5–3.5 mm. broad, obtuse or rotund and pubescent at the apex, otherwise glabrous; ligulate flowers 8, tube slender, 3–3.5 mm. long, limb usually radiate or sometimes irregularly dentate and occasionally bearing reduced stamens, yellow; flowers of the disk 25–30 with a tubular-campanulate, equally 5-lobed, yellow corolla; setae of the pappus copious, white; achenes glabrous.

Colombia: in wet places, foot of Cordillera Oriental, near Neiva, August 1, 1917, *Rusby & Pennell 572* (N. Y. Bot. Gard. Herb., fragment and photograph in Mo. Bot. Gard. Herb.), TYPE.

Specimens apparently representing this species, but with somewhat smaller heads, and intermediate between it and the following variety, were collected along the Camino de Gachetá, Colombia, January 5, 1920, *Bro. Ariste-Joseph 467 A* (U. S. Nat. Herb.).

Var. *gachetensis* Greenm., var. nov.

Stem arachnoid-tomentulose intermixed with multicellular somewhat villous hairs; leaves about 10 cm. long, one-half as broad, persistently subarachnoid-tomentulose beneath; heads about 20-flowered.

Colombia: Camino de Gachetá, January, 1920, *Bro. Ariste-Joseph A 535* (U. S. Nat. Herb., No. 1059490).

This species is related apparently to *S. Urbani* Hieron. but differs in having larger leaves, longer petioles, shorter involucre bracts, and a distinctly white pappus.

Senecio pericaulis Greenm., sp. nov.

Perennis; caulibus verisimiliter scandentibus lignescentibus teretibus striatis brunneis hirtellis plus minusve glabris; foliis alternis sessilibus amplexicaulibusque anguste ovatis vel oblongo-ovatis 4–8 cm. longis 1.5–3 cm. latis mucronato-acutis callosodenticulatis vel leviter sinuato-dentatis utrinque aliquantulum crispo-hirtellis plus minusve glabris, marginibus saepe revolutis; inflorescentiis terminalibus laxo paniculato-cymosis subglabris vel sparse glanduloso-hirtellis; pedunculis 1.5–4.5 cm. longis; capitulis 12–14 mm. altis radiatis; involucri campanulatis, calyculatis; bracteis involucri 13 lineari-lanceolatis vel lanceolato-oblongis circiter 1 cm. longis 1–3 mm. latis acutis vel obtusis glabris vel sparsissime glanduloso-hirtellis; floribus liguliferis 10–13, tubo 5 mm. longo, ligula oblonga circiter 12 mm. longa 3–4 mm. lata pallide aurantia; floribus disci circiter 30, corollis 8 mm. longis sursum sensim ampliatis; pappi setis albidis; achaeniis glabris.

Perennial; stems probably scandent, ligneous, terete, striate, brownish, crisp-hirtellous, more or less glabrate; leaves alternate, sessile and amplexicaul, narrowly ovate or oblong-ovate, 4–8 cm. long, 1.5–3 cm. broad, mucronate-acute, callous-denticulate or shallowly sinuate-dentate, somewhat crisp-hirtellous on both surfaces, more or less glabrate, especially above, margins commonly revolute; inflorescence a terminal subglabrous or sparing-

ly glandular-hirtellous open paniculate cyme; peduncles 1.5-4.5 cm. long; heads 12-14 mm. high, radiate; involucre campanulate, calyculate with few short setaceous bracteoles; bracts of the involucre 13, linear-lanceolate or lanceolate-oblong, about 1 cm. long, 1-3 mm. broad, acute or obtuse, glabrous or very sparingly glandular-hirtellous; ray-flowers 10-13, tube 5 mm. long, ligule oblong, about 1 cm. long, 3-4 mm. broad, pale yellow; disk-flowers about 30, corollas gradually amplified from the base; setae of the pappus white; achenes glabrous.

Ecuador: mountains, Province of Cuenca, July, 1864, *Jameson*, without number (U. S. Nat. Herb., No. 700136), TYPE; vicinity of Cumbe, September 25, 1918, *Rose, Pachano & Rose 22983* (U. S. Nat. Herb., No. 1023355).

This species is allied to *S. amplexifolius* HBK., from which it differs in having smaller and more or less crisp-hirtellous leaves, a somewhat glandular-hirtellous inflorescence, and fewer flowers in the head.

Senecio pindilicensis Hieron. in Engl. Bot. Jahrb. 19:65. 1894.

Ecuador: Quito, without number or date of collection, *Jameson* (Gray Herb.).

The specimen here cited agrees in all essential details with the very careful description of the above species by Professor Hieronymus, and the writer has no hesitation in referring it to that species.

Senecio pulchellus DC. Prodr. 6:421. 1837; Wedd. Chlor. And. 1:100. 1855-1857.

Cacalia pulchella HBK. Nov. Gen. & Sp. 4:160. 1820.

Microchaete pulchella Benth. Pl. Hartw. 210. 1845.

M. trichopus Benth. Pl. Hartw. 210. 1845.

Colombia: vicinity of Bogotá, alt. 2895 m., *Hartweg 1162* (N. Y. Bot. Gard. Herb.); in woods between Huambia and Pitaya, Province of Popaya, *Hartweg 1163* (N. Y. Bot. Gard. Herb.); dry páramo, southeast of Guadalupe, near Bogotá, Department of Cundinamarca, October 12, 1917, *Pennell 2377* (N. Y. Bot. Gard. Herb., Mo. Bot. Gard. Herb., and Gray Herb.).

The leaves on Dr. Pennell's specimen here cited are somewhat larger than described by Kunth for *Cacalia pulchella* on which the above species was founded, but in all other regards there is complete accord with that species.

Senecio rudbeckiaefolius Meyen & Walp. in Nov. Act. Nat. Cur. 19: Suppl. 1, p. 283. 1843; Walp. Rep. 6:272. 1846-47.

Bolivia: Rio Tapacari, March 19, 1892, *Kuntze* (U. S. Nat. Herb.); Cochabamba, February 28, 1920, *Holway & Holway* 343 (U. S. Nat. Herb.).

***Senecio sanctae-martae* Greenm., sp. nov.**

Caulis lignescens verisimiliter scandens; ramis floriferis terebibus striatis brunneis glabris; foliis oppositis sessilibus late ovatis 5-7 cm. longis 3-4.5 cm. latis crenato-serratis vel supremis subintegerrimis acutis vel obtusis basi cordatis amplexicaulibusque utrinque glabris et plus minusve glaucescentibus; inflorescentiis terminalibus corymboso-cymosis parce pubescentibus multicapitatis; capitulis 1 cm. altis heterogamis; involucri anguste campanulatis brevis calyculatis; involucri squamis plerumque 13 lineari-lanceolatis 6-7 mm. longis acutis glabris; ligulis anguste oblongis circiter 1 cm. longis flavis; floribus disci 30-35; pappi setis albis; achaeniis glabris.

Stem ligneous, probably scandent; branches terete, striate, glabrous; leaves opposite, sessile, broadly ovate, 5-7 cm. long, 3-4.5 cm. broad, crenate-serrate or the uppermost entire or nearly so, acute or obtuse at the apex, cordate and amplexicaul at the base, glabrous on both surfaces and more or less glaucous; inflorescence a terminal round-topped many-headed corymbose cyme, sparingly pubescent; heads about 1 cm. high, radiate; involucre narrowly campanulate, calyculate with short bracteoles; bracts of the involucre usually 13, linear-lanceolate, 6-7 mm. long, acute, glabrous; ray-flowers 8-10, rays narrowly oblong, about 1 cm. long, yellow; disk-flowers 30-35; pappus white; achenes glabrous.

Colombia: Santa Marta, Province of Magdalena, 1898-1899, *Herbert H. Smith* 1977 (Mo. Bot. Gard. Herb., Field Mus. Nat. Hist. Herb., and N. Y. Bot. Gard. Herb.), TYPE.

Although no notes are recorded concerning the plant here described, yet the specimens at hand indicate a more or less scandent habit. The species, however, is very characteristic on account of the opposite, sessile, glabrous, and somewhat glaucous leaves.

***Senecio scortifolius* Greenm., sp. nov.**

Perennis; caulibus verisimiliter scandentibus lignescentibus brunneis striatis glabris; foliis alternis petiolatis ovato-ellipticis 3-11 cm. longis 1.5-6 cm. latis acutis vel obtusis vel raro subrotundatis integris basi cuneatis utrinque glabris coriaceis et in sicco saepe pallido-viridibus; petioliis .5-2 cm. longis; inflorescentiis in speciminibus axillaribus paniculatis glabris vel paululo

DARTMOUTH
COLLEGE
LIBRARY

puberulentis, pedunculis usque ad 12 cm. longis glabris; capitulis numerosis 7-8 mm. latis radiatis; involucris campanulatis parce calyculatis; involucris bracteis 8 lineari-lanceolatis vel lanceolato-oblongis 4.5-5 mm. longis acutis vel obtusis ad apicem penicillatis ceterum glabris; floribus liguliferis 3-5, tubulo circiter 3 mm. longo, ligula anguste oblonga 3.5 mm. longa flava; floribus disci 11-15, tubulo in limbum sensim ampliatus transeunte; pappi setis albidis usque ad 6 mm. longis; achaeniis glabris.

Perennial; stems probably scandent, ligneous, brownish, striate, glabrous; leaves alternate, petiolate, ovate-elliptic, 3-11 cm. long, 1.5-6 cm. broad, acute or obtuse, or rarely subrotund, entire, cuneate at the base, glabrous on both surfaces, coriaceous, and in the dried state usually pale or dull green; petioles .5-2 cm. long; inflorescence in specimens at hand in axillary glabrous or sparingly puberulent panicles, peduncles as much as 12 cm. in length, glabrous; heads numerous, 7-8 mm. high, radiate; involucre campanulate, sparingly calyculate; bracts of the involucre 8, linear-lanceolate or lanceolate-oblong, 4.5-5 mm. long, acute or obtuse, penicillate at the apex, otherwise glabrous; ligulate flowers 3-5, tube about 3 mm. long, ligule narrowly oblong, 3.5 mm. long, yellow; flowers of the disk 11-15, tube gradually amplified into the limb; setae of the pappus white, 5-6 mm. long; achenes glabrous.

Colombia: Santa Marta, alt. 1400 m., coll. of 1898-1901, *Herbert H. Smith 2000* (Mo. Bot. Gard. Herb., TYPE; Gray Herb., Phil. Acad. Nat. Sci. Herb., U. S. Nat. Herb., and Field Mus. Nat. Hist. Herb.).

Venezuela: Páramo de Aricagua, Mérida, alt. 2500 m., March 31, 1922, *Jahn 1032* (U. S. Nat. Herb.).

The species here described has many characters in common with *S. ellipticifolius* Hieron., but that species is described as a shrub with glandular puberulent peduncles and bracteoles, and with somewhat shining leaves. *S. scortifolius* from the several specimens at hand presents every appearance of a scandent plant, and the leaves in the dried state are dull green and are doubtless rather fleshy in the living state. In foliar characters *S. scortifolius* suggests *S. epiphyticus* Kuntze of Bolivia, which has a pronounced fuscous pubescence in the inflorescence and on the involucre. It also simulates *S. Tonduzii* Greenm. of Costa Rica.

Senecio sinapoides Rusby in Mem. Torr. Bot. Club 6:65. 1896.

Bolivia: vicinity of Cochabamba, coll. of 1891, *Bang 1135*

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS
VOLUME 1135

(Mo. Bot. Gard. Herb., U. S. Nat. Herb., and Phil. Acad. Nat. Sci. Herb.); Cochabamba, March 5, 1920, *Holway & Holway 371* (Gray Herb. and U. S. Nat. Herb.).

Senecio Sprucei Klatt in *Leopoldina* 24:128. 1887.

Peru: near Tarapota, coll. of 1855-56, *Spruce 3926* (Gray Herb.), CO-TYPE.

Senecio stigophlebius Baker in *Mart. Fl. Bras.* 6³:321. 1884.

Brazil: vicinity of Itatiaya, July 26-30, 1915, *Rose & Russell 20504* and *20549* (U. S. Nat. Herb., fragment and photograph in Mo. Bot. Gard. Herb.).

The plant of Rose and Russell accords with Sello's No. 2187, a specimen of which is in the Gray Herbarium, which is the type-number of the above species.

Senecio Stuebelii Hieron. in *Engl. Bot. Jahrb.* 21:357. 1895.

Colombia: wet mossy páramo, Páramo de Cruz Verde, near Bogotá, Department of Cundinamarca, alt. 3300-3400 m., September 20, 1917, *Pennell 2055* (N. Y. Bot. Gard. Herb., Gray Herb., U. S. Nat. Herb., and Mo. Bot. Gard. Herb.); moist páramo, southeast of Guadalupe, near Bogotá, alt. 3300-3400 m., September 27, 1917, *Pennell 2255* (N. Y. Bot. Gard. Herb., U. S. Nat. Herb., Gray Herb., Field Mus. Nat. Hist. Herb., and Mo. Bot. Gard. Herb.); Páramo de Choachi, near Bogotá, alt. 3700 m., August 8, 1922, *Killip & Bro. Ariste-Joseph 11975* (U. S. Nat. Herb.).

Senecio subdecurrens Schz. Bip. in *Bonplandia* 4:55. 1856, name only; Wedd. *Chlor. And.* 1:109. 1855-57, description.

S. formosus Rusby in *Bull. N. Y. Bot. Gard.* 4:393. 1907, not HBK.

Colombia: Páramo de Buena Vista, Huila group, Central Cordillera, State of Cauca, alt. 3000-3600 m., January, 1906, *Pittier 1130* (U. S. Nat. Herb.).

Ecuador: without number, coll. of December, 1876, *Andre* (Gray Herb.); also *Spruce 6001* (Gray Herb.).

Bolivia: without locality or date of collection, *Bang 1958* (Mo. Bot. Gard. Herb. and Phil. Acad. Nat. Sci. Herb.).

Mr. Bang's plant was distributed as "*Senecio formosus* HBK.," under which name it may be found in herbaria.

Senecio subglomerosus Greenm., sp. nov.

Plate 7.

Frutex scandens usque ad 3 m. altus; caulibus ramisque terebibus striatis fulvo-tomentulosus; foliis alternis petiolatis lanceolatis vel elliptico-lanceolatis 3-10 cm. longis 1.5-3 cm. latis saepe acuminatis acutis integris juventate parce fusco-puberu-

lentis plus minusve glabratis pallido-viridibus, medio nervio venisque immersis et supra persistenter puberulentis subtus aliquanto prominentibus; inflorescentiis terminalibus paniculatis fusco-tomentulosis; capitulis subdiscoideis sed heterogamis circiter 8 mm. altis in glomerulis dispositis; involucris anguste campanulatis calyculatis; involucris bracteis 8-13 lineari-lanceolatis 5-7 mm. longis acutis penicillatis extrinsecus fusco-puberulentis; floribus femineis 3-5 multo reductis, corollis nonligulatis gracilibus tubuliformibus circiter 4 mm. longis subaequaliter 5-dentatis; floribus disci 10-12 hermaphroditis, corollis circiter 6 mm. longis tubulo-campanulatis aequaliter 5-dentatis; pappi setis albidis; achaeniis glabris vel sparsissime hirtellis.

Scandent shrub, 3 m. or less in height; stem and branches terete, striate, tawny tomentulose; leaves alternate, petiolate, lanceolate or elliptic-lanceolate, 3-10 cm. long, 1.5-3 cm. broad, usually acuminate, acute, entire, sparingly tawny puberulent in the early stages, more or less glabrate, pale green, the median nerve and veins immersed and persistently puberulent above, somewhat prominent beneath; inflorescence a terminal tawny tomentulose panicle; heads subdiscoid but heterogamous, about 8 mm. high, disposed in glomerules; involucre narrowly campanulate, calyculate; bracts of the involucre 8-13, linear-lanceolate, 5-7 mm. long, acute, penicillate, externally tawny puberulent; female flowers 3-5, much reduced, corollas nonligulate, slender, tubuliform, about 4 mm. long, subequally 5-dentate; flowers of the disk 10-12, hermaphrodite, corollas tubular-campanulate, about 6 mm. long, equally 5-dentate; setae of the pappus white; achenes glabrous or very sparingly hirtellous.

Bolivia: Coroico, Yungas, September, 1894, *Bang* 2459, (N. Y. Bot. Gard. Herb., TYPE; U. S. Nat. Herb., Gray Herb., and Mo. Bot. Gard. Herb.).

The relationship of this species seems to be with *S. Jelskii* Hieron. from which, however, it differs in having persistent tomentose branches, tawny puberulent leaves, and rayless pistillate flowers.

***Senecio subbruncinatus* Greenm., comb. nov.**

S. formosus var. *subbruncinatus* Wedd. Chlor. And. 1:125. 1855-57.

Herba perennis; caule erecto vel adscendente simplici vel superne ramoso .5-1 m. alto glabro vel sparsissime crispo-puberulento striato plus minusve purpurascenti; foliis inferioribus oblanceolatis 5-10 cm. longis 1-1.5 cm. latis obtusis vel submu-

cronato-acutis sinuato-dentatis basi in petiolo contractis, superioribus sessilibus plus minusve amplexicaulibus lanceolatis 2-12 cm. longis usque ad 1.5 cm. latis grosse dentatis vel subbruncinatis glabris vel subglabris, eis ad inflorescentias plerumque multo reductis; inflorescentiis terminalibus subcorymboso-cymosis paucicapitatis vel aliquando unicapitatis; capitulis magnis 15-18 mm. altis radiatis; involucris campanulatis calyculatis; involucri bracteis 21 lanceolatis 10-13 mm. longis 1.5-3 mm. latis acutis penicillatis glabris vel parcissime crispo-puberulentis purpurascens; floribus ligulatis circiter 21, tubulo 5 mm. longo, ligula anguste oblonga circiter 12 mm. longa 2-3 mm. lata violaceo-purpurea; floribus disci numerosis; pappi setis albidis; achaeniis parcissime hirtellis.

Perennial herb; stem erect or ascending, simple or branched above, .5-1 m. high, glabrous or sparingly crisp-puberulent, striate, more or less purple; lower leaves oblanceolate, 5-10 cm. long, 1-1.5 cm. broad, obtuse or submucronate-acute, sinuate-dentate, contracted at the base into a narrowly winged petiole; upper leaves sessile, more or less amplexicaul, lanceolate, 2-12 cm. long, 1.5 cm. or less broad, coarsely dentate to subbruncinate, glabrous or nearly so, those towards the inflorescence usually much reduced; inflorescence a terminal subcorymbose cyme, few-headed or occasionally one-headed; heads large, 15-18 mm. high, radiate; involucre campanulate, calyculate; bracts of the involucre 21, lanceolate, 10-13 mm. long, 1.5-3 mm. broad, acute, penicillate, glabrous or sparingly crisp-puberulent, purple; ligulate flowers about 21, tube 5 mm. long, ligule narrowly oblong, about 12 mm. long, 2-3 mm. broad, violet-purple; flowers of the disk numerous, more than 100; setae of the pappus white; achenes sparingly hirtellous.

Colombia: wet mossy páramo, Páramo de Cruz Verde, near Bogotá, Department of Cundinamarca, alt. 3300-3500 m., September 20, 1917, *Pennell 2051* (N. Y. Bot. Gard. Herb., U. S. Nat. Herb., and Mo. Bot. Gard. Herb.); Canoas, July, 1917, *Bro. Ariste-Joseph A 87* (U. S. Nat. Herb.).

The writer feels confident that this is the form which Weddell briefly characterized as *Senecio formosus* var. *subbruncinatus*; but the slender lower part of the stem, the oblanceolate to subbruncinate leaves, and the glabrous or essentially glabrous character of the entire plant seems to warrant its specific rank.

Senecio sylvicolus Greenm., sp. nov.

Plate 8.

Frutex scandens; ramis ramulisque brunneis striatis glabris;

foliis alternis petiolatis elliptico-oblongis vel ovatis basi cuneatis apice submucronatis integris utrinque glabris supra fusco-viridibus subtus pallidioribus, nervo medio et nervis lateralibus prominentibus; petiolis 5-10 mm. longis canaliculatis glabris; inflorescentiis terminalibus corymboso-cymosis parce crispo-pubescentibus; capitulis pedunculatis heterogamis circiter 12 mm. altis; involucris campanulatis parce calyculatis floccoso-tomentulosis; involucri bracteis 8 lanceolatis vel lanceolato-oblongis circiter 6 mm. longis 1-2 mm. latis obtusis penicillatisque; floribus liguliferis 5, ligulis anguste oblongis 10-12 mm. longis 5-nervatis flavibus; floribus disci 12-15; pappi setis albidis; achaeniis glabris.

Stem scandent; branches and branchlets brownish, terete, striate, glabrous; leaves alternate, petiolate, elliptic-oblong or ovate, cuneate at the base, submucronate, entire, glabrous on both surfaces, dark green above, paler beneath, midrib and lateral nerves prominent; petioles 5-10 mm. long, canaliculate, glabrous; inflorescence terminating the stem and lateral branches in sparingly crisp-pubescent corymbose cymes; heads pedunculate, about 12 mm. high; involucre campanulate, sparingly calyculate with minute bracteoles, floccose-tomentulose; bracts of the involucre 8, lanceolate or lanceolate-oblong, about 6 mm. long, 1-2 mm. broad, obtuse, penicillate; ray-flowers 5, rays narrowly oblong, 10-12 mm. long, 5-nerved, yellow; disk-flowers 12-15; pappus white; achenes glabrous.

Colombia: edge of forest, "Rosalito," near Páramo de Ruiz, Department of Tolima, alt. 2800-3100 m., December 15-17, 1917, *Pennell 2985* (N. Y. Bot. Gard. Herb., Mo. Bot. Gard. Herb., Gray Herb., U. S. Nat. Herb., and Field Mus. Nat. Hist. Herb.).

This species is related to *S. ellipticus* DC. and *S. ellipticifolius* Hieron. From the former it differs in having smaller leaves and a tomentulose inflorescence; and from the latter it differs in the absence of glandular puberulence, longer petioles, and in the absence of conspicuous calyculiform bracts subtending the involucre.

***Senecio teretifolius* (HBK.) DC. Prodr. 6:420. 1837.**

Cacalia teretifolia HBK. Nov. Gen. & Sp. 4:159, pl. 357. 1820.

Ecuador: Palmira, alt. 3200 m., August 23, 1918, *Rose & Rose 22311* (U. S. Nat. Herb. and N. Y. Bot. Gard. Herb.); vicinity of Ambato, Province of Tungurahua, December, 1918, *Pachano 19* (U. S. Nat. Herb. and N. Y. Bot. Gard. Herb.); vicinity of Ambato, August 24-26, 1918, *Rose & Rose 22327* (U. S. Nat. Herb.).

Senecio theaeifolius Benth. Pl. Hartw. 210. 1845; Walp. Rep. 6:271. 1846-47.

Ecuador: without further data, *Jameson* (U. S. Nat. Herb. No. 534598).

Bolivia: near Bogotá, *Hartweg 1166* (Gray Herb.).

Senecio tolimensis Schz. Bip. ex Wedd. Chlor. And. 1:98. 1855-57.

Colombia: sphagnum bog, "Balsillas," on Rio Balsillas, Department of Huila, alt. 2100-2200 m., August 3-5, 1917, *Rusby & Pennell 765* (N. Y. Bot. Gard. Herb. and Mo. Bot. Gard. Herb.); near Bogotá, colls. of 1919 and 1920, *Bro. Ariste-Joseph*, without number (U. S. Nat. Herb.); Páramo de Guasca, January 3, 1920, *Bro. Ariste-Joseph A 478* (U. S. Nat. Herb. and Gray Herb.).

Senecio vaccinioides Schz. Bip. ex Wedd. Chlor. And. 1:99 pl. 20, fig. A. 1855-57.

Cacalia vaccinioides HBK. Nov. Gen. & Sp. 4:161, pl. 358. 1820.

Psacalium vaccinioides DC. Prodr. 6:335. 1837.

Microchaete vaccinioides Benth. Pl. Hartw. 210. 1845.

Colombia: Páramo de Buena Vista, Huila group, Central Cordillera, State of Cauca, alt. 3000-3600 m., January, 1906, *Pittier 1181* (U. S. Nat. Herb.); dry grassy páramo, Cerro de Focha, near Bogotá, Department of Cundinamarca, alt. 3100-3200 m., December 26, 1917, *Pennell 2198* (N. Y. Bot. Gard. Herb.); moist open places, Páramo de Ruiz, Department of Tolima, alt. 3400-3700 m., December 16, 17, 1917, *Pennell 3073* (N. Y. Bot. Gard. Herb.).

Var. **pruinosis** Wedd. Chlor. And. 1:99. 1855-57.

Cacalia glabrata HBK. Nov. Gen. & Sp. 4:161. 1820.

Psacalium glabratum DC. Prodr. 6:335. 1837.

Microchaete glabrata Benth. Pl. Hartw. 210. 1845.

Colombia: grassy páramo, Páramo de Chaquiro, Cordillera Occidental, Department of Bolivar-Antioquia, alt. 3000-3200 m., February 23, 1918. *Pennell 4261* (N. Y. Bot. Gard. Herb.); Alto del Verjón, *Bro. Ariste-Joseph 1-A* (U. S. Nat. Herb.).

Senecio Weddellianus Hieron. in Engl. Bot. Jahrb. 21:356. 1895.

S. vernicosus Schz. Bip. var. *microphyllus* Wedd. Chlor. And. 1:94. 1855-1857.

Colombia: moist open places, Páramo de Ruiz, Department of Tolima, alt. 3500-3800 m., December 16-17, 1917, *Pennell*

3002 (N. Y. Bot. Gard. Herb., U. S. Nat. Herb., and Mo. Bot. Gard. Herb.).

Senecio Xanthopappus Klatt ex Hieron. in Engl. Bot. Jahrb. 21:360. 1895.

Colombia: roadside, Sibate, Department of Cundinamarca, October 13-15, 1917, *Pennell 2514* (N. Y. Bot. Gard. Herb., fragment and photograph in Mo. Bot. Gard. Herb.); Bogotá, coll. of 1917, *Bro. Ariste-Joseph A 9* (U. S. Nat. Herb.); Sabana de Bogotá, May, 1923, *G. H. Pring* (Mo. Bot. Gard. Herb.); San Cristóbal, October, 1918, *Bro. Ariste-Joseph A 267* (U. S. Nat. Herb., Mo. Bot. Gard. Herb., N. Y. Bot. Gard. Herb., and Gray Herb.).

EXPLANATION OF PLATE

PLATE 3


Senecio aberrans Greenm.

Colombia

From the type specimen, Pennell No. 3636, in the New York Botanical Garden Herbarium.



GREENMAN—SOUTH AMERICAN SENECIOS




EXPLANATION OF PLATE

PLATE 4

Senecio aridus Greenm.

Colombia

From the type specimen, Pennell No. 3037, in the New York Botanical Garden Herbarium.



GREENMAN—SOUTH AMERICAN SENECIOS

EXPLANATION OF PLATE

PLATE 5


Senecio lophophilus Greenm.


Colombia

From the type specimen, Herbert H. Smith, No. 1988, in the New York Botanical Garden Herbarium.



GREENMAN—SOUTH AMERICAN SENECIOS





EXPLANATION OF PLATE

PLATE 6


Senecio Pennellii Greenm.

Colombia

From the type specimen, Rusby & Pennell No. 572, in the New York Botanical Garden Herbarium.



GREENMAN—SOUTH AMERICAN SENECIOS



EXPLANATION OF PLATE

PLATE 7

Senecio subglomeratus Greenm.

Bolivia

From the type specimen, Bang No. 2459, in the New York Botanical Garden Herbarium.






EXPLANATION OF PLATE

PLATE 8

Senecio sylvicolus Greenm.

Colombia

From the type specimen, Pennell No. 2985, in the New York Botanical Garden Herbarium.



GREENMAN—SOUTH AMERICAN SENECIOS



